

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 15/18, 15/56, 15/58, 5/00, 9/04

(52) International Filing Date: 11 January 1997 (17.01.97)

(53) International Application Number: PCT/US97/03593

(54) Designated States: CA, JP, EA, RU, BY, AZ, BE, CH, DE, DK, ES, FI, FR, GR, GB, IE, IT, LU, MC, NL, PT, SE, SI

(55) International Publication Date: 24 July 1997 (24.07.97)

(56) Priority Date: 08.08.95

(57) Applicant: AVIDON, INC. (US); Suite 1200, 1701 Harbor Blvd., Alhambra, CA 91801 (US); AVIDON CORPORATION, UNIVERSITY (US); 720 Ruben Avenue, Baltimore, MD 21203 (US)

(58) Inventor: PODSACOF, Gregory M.; 2932 Ted Drive, Pasadena, CA 91633 (US); NEISLER, Paul D.; 7932 Eubank Avenue, Baltimore, MD 21234 (US); BYRNE, Barry J.; 814 Cummer Road, Baltimore, MD 21113 (US); KORTZMAN, Gary J.; 24 Cuneel Lane, Middle Park, CA, 94023 (US)

(59) Agent: MCKRACKEN, Thomas P. & J.; Reine & Associates, Suite 200, 20 Midway Road, Middle Park, CA 94023 (US)

(60) Title: METHODS FOR DELIVERING DNA TO MUSCLE CELLS USING RECOMBINANT ADENOVIRUS ASSOCIATED VIRUS

(61) Abstract

The use of recombinant adenovirus (AAV) vectors for delivery of DNA molecules to muscle cells and tissue is disclosed. A series of in vivo transductions of muscle cells with an adenovirus AAV vector have been made. These cells, by intramuscular injection, as well as oral gavage, have been shown to be capable of expressing the delivered gene and for in vivo secretion of the expressed protein from transduced muscle cells.

(62) Summary of the Invention

The use of recombinant adenovirus (AAV) vectors for delivery of DNA molecules to muscle cells and tissue is disclosed. A series of in vivo transductions of muscle cells with an adenovirus AAV vector have been made. These cells, by intramuscular injection, as well as oral gavage, have been shown to be capable of expressing the delivered gene and for in vivo secretion of the expressed protein from transduced muscle cells.

(63) Claims

1. A method for delivering a DNA molecule to muscle cells and tissue, comprising: (a) providing a recombinant adenovirus (AAV) vector containing a DNA molecule; (b) administering the AAV vector to a subject; and (c) detecting the presence of the DNA molecule in the muscle cells and tissue of the subject.

2. The method of claim 1, wherein the DNA molecule is a gene.

3. The method of claim 1, wherein the DNA molecule is a protein-coding sequence.

4. The method of claim 1, wherein the DNA molecule is a non-coding sequence.

5. The method of claim 1, wherein the DNA molecule is a gene encoding a protein.

6. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

7. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

8. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

9. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

10. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

11. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

12. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

13. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

14. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

15. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

16. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

17. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

18. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

19. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

20. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

21. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

22. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

23. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

24. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

25. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

26. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

27. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

28. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is secreted from the muscle cells and tissue.

29. The method of claim 1, wherein the DNA molecule is a gene encoding a protein that is expressed in the muscle cells and tissue.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front page of pamphlets (symbolizing international application) under the PCT.

AF	Algeria	GA	Ghana	MY	Malaysia
AG	Argentina	GB	Great Britain	MX	Mexico
AI	Angola	GR	Greece	NI	Nicaragua
AL	Albania	HR	Croatia	NO	Norway
AM	Armenia	HU	Hungary	PL	Poland
AN	Antigua and Barbuda	IE	Ireland	PT	Portugal
AO	Angola	IL	Israel	RO	Romania
AP	Andorra	IN	India	RU	Russian Federation
AR	Argentina	JP	Japan	SA	Saudi Arabia
AS	Armenia	KE	Kenya	SE	Sweden
AT	Austria	KR	Republic of Korea	SG	Singapore
AU	Australia	KZ	Kazakhstan	SI	Slovenia
AW	Aruba	LA	Laos	SK	Slovakia
AX	Aland Islands	LB	Lebanon	SN	Senegal
AY	Ascension Island	LC	Liechtenstein	SO	Somalia
AZ	Azerbaijan	LI	Liechtenstein	SR	Suriname
BA	Bosnia and Herzegovina	LK	Sri Lanka	SV	El Salvador
BB	Barbados	LU	Luxembourg	TD	Chad
BC	Bahamas	LV	Latvia	TE	Togo
BD	Bangladesh	LY	Libya	TH	Thailand
BE	Belgium	MA	Morocco	TI	Timor-Leste
BF	Burkina Faso	MC	Monaco	TM	Turkmenistan
BG	Bulgaria	MD	Moldova	TN	Tunisia
BH	Bahrain	ME	Montenegro	TR	Turkey
BI	Burundi	MG	Madagascar	UA	Ukraine
BJ	Benin	MH	Marshall Islands	UG	Uganda
BK	Belize	MI	Maldives	US	United States of America
BL	Bolivia	MM	Myanmar	UY	Uruguay
BM	Bermuda	NA	Namibia	UZ	Uzbekistan
BN	Brunei Darussalam	NE	Niger	VE	Venezuela
BO	Bolivia	NG	Nigeria		
BR	Brazil	NL	Netherlands		
BS	Bahamas	NI	Nicaragua		
BT	Bhutan	NL	Netherlands		
BV	Bouvet Island	NO	Norway		
BW	Botswana	NP	Nepal		
BY	Belarus	NZ	New Zealand		
BZ	Belize	OM	Oman		
CA	Canada	PA	Panama		
CB	Cambodia	PE	Peru		
CC	Cocos (Keeling) Islands	PG	Papua New Guinea		
CD	Congo	PH	Philippines		
CE	Cote d'Ivoire	PK	Pakistan		
CF	Cote d'Ivoire	PL	Poland		
CG	Congo	PT	Portugal		
CH	Switzerland	RO	Romania		
CI	Cote d'Ivoire	RU	Russian Federation		
CJ	Czech Republic	SA	Saudi Arabia		
CK	Cook Islands	SE	Sweden		
CL	Chile	SG	Singapore		
CM	Cameroun	SI	Slovenia		
CN	China	SK	Slovakia		
CO	Colombia	SN	Senegal		
CR	Costa Rica	SO	Somalia		
CU	Cuba	SR	Suriname		
CV	Cape Verde	SV	El Salvador		
CW	Curaçao	TD	Chad		
CX	Christmas Island	TE	Togo		
CY	Cyprus	TH	Thailand		
CZ	Czech Republic	TI	Timor-Leste		
DA	Denmark	TM	Turkmenistan		
DB	Dominican Republic	TN	Tunisia		
DE	Germany	TR	Turkey		
DF	Dominican Republic	UA	Ukraine		
DG	Dominican Republic	UG	Uganda		
DH	Dominican Republic	US	United States of America		
DI	Dominican Republic	UY	Uruguay		
DJ	Dominican Republic	UZ	Uzbekistan		
DK	Denmark	VE	Venezuela		
DL	Dominican Republic				
DM	Dominican Republic				
DN	Dominican Republic				
DO	Dominican Republic				
DP	Dominican Republic				
DQ	Dominican Republic				
DR	Dominican Republic				
DS	Dominican Republic				
DT	Dominican Republic				
DU	Dominican Republic				
DV	Dominican Republic				
DW	Dominican Republic				
DX	Dominican Republic				
DY	Dominican Republic				
DZ	Dominican Republic				
EA	Dominican Republic				
EB	Dominican Republic				
EC	Dominican Republic				
ED	Dominican Republic				
EE	Dominican Republic				
EF	Dominican Republic				
EG	Dominican Republic				
EH	Dominican Republic				
EI	Dominican Republic				
EJ	Dominican Republic				
EK	Dominican Republic				
EL	Dominican Republic				
EM	Dominican Republic				
EN	Dominican Republic				
EO	Dominican Republic				
EP	Dominican Republic				
EQ	Dominican Republic				
ER	Dominican Republic				
ES	Dominican Republic				
ET	Dominican Republic				
EU	Dominican Republic				
EV	Dominican Republic				
EW	Dominican Republic				
EX	Dominican Republic				
EY	Dominican Republic				
EZ	Dominican Republic				
FA	Dominican Republic				
FB	Dominican Republic				
FC	Dominican Republic				
FD	Dominican Republic				
FE	Dominican Republic				
FF	Dominican Republic				
FG	Dominican Republic				
FH	Dominican Republic				
FI	Dominican Republic				
FJ	Dominican Republic				
FK	Dominican Republic				
FL	Dominican Republic				
FM	Dominican Republic				
FN	Dominican Republic				
FO	Dominican Republic				
FP	Dominican Republic				
FQ	Dominican Republic				
FR	Dominican Republic				
FS	Dominican Republic				
FT	Dominican Republic				
FU	Dominican Republic				
FV	Dominican Republic				
FW	Dominican Republic				
FX	Dominican Republic				
FY	Dominican Republic				
FZ	Dominican Republic				
GA	Dominican Republic				
GB	Dominican Republic				
GC	Dominican Republic				
GD	Dominican Republic				
GE	Dominican Republic				
GF	Dominican Republic				
GG	Dominican Republic				
GH	Dominican Republic				
GI	Dominican Republic				
GJ	Dominican Republic				
GK	Dominican Republic				
GL	Dominican Republic				
GM	Dominican Republic				
GN	Dominican Republic				
GO	Dominican Republic				
GP	Dominican Republic				
GQ	Dominican Republic				
GR	Dominican Republic				
GS	Dominican Republic				
GT	Dominican Republic				
GU	Dominican Republic				
GV	Dominican Republic				
GW	Dominican Republic				
GX	Dominican Republic				
GY	Dominican Republic				
HA	Dominican Republic				
HB	Dominican Republic				
HC	Dominican Republic				
HD	Dominican Republic				
HE	Dominican Republic				
HF	Dominican Republic				
HG	Dominican Republic				
HH	Dominican Republic				
HI	Dominican Republic				
HJ	Dominican Republic				
HK	Dominican Republic				
HL	Dominican Republic				
HM	Dominican Republic				
HN	Dominican Republic				
HO	Dominican Republic				
HP	Dominican Republic				
HQ	Dominican Republic				
HR	Dominican Republic				
HS	Dominican Republic				
HT	Dominican Republic				
HU	Dominican Republic				
HV	Dominican Republic				
HW	Dominican Republic				
HX	Dominican Republic				
HY	Dominican Republic				
HZ	Dominican Republic				
IA	Dominican Republic				
IB	Dominican Republic				
IC	Dominican Republic				
ID	Dominican Republic				
IE	Dominican Republic				
IF	Dominican Republic				
IG	Dominican Republic				
IH	Dominican Republic				
II	Dominican Republic				
IJ	Dominican Republic				
IK	Dominican Republic				
IL	Dominican Republic				
IM	Dominican Republic				
IN	Dominican Republic				
IO	Dominican Republic				
IP	Dominican Republic				
IQ	Dominican Republic				
IR	Dominican Republic				
IS	Dominican Republic				
IT	Dominican Republic				
IU	Dominican Republic				
IV	Dominican Republic				
IW	Dominican Republic				
IX	Dominican Republic				
IY	Dominican Republic				
IZ	Dominican Republic				
JA	Dominican Republic				
JB	Dominican Republic				
JC	Dominican Republic				
JD	Dominican Republic				
JE	Dominican Republic				
JF	Dominican Republic				
JG	Dominican Republic				
JH	Dominican Republic				
JI	Dominican Republic				
JJ	Dominican Republic				
JK	Dominican Republic				
JL	Dominican Republic				
JM	Dominican Republic				
JN	Dominican Republic				
JO	Dominican Republic				
JP	Dominican Republic				
JQ	Dominican Republic				
JR	Dominican Republic				
JS	Dominican Republic				
JT	Dominican Republic				
JU	Dominican Republic				
JV	Dominican Republic			</	

5  
METHODS FOR DELIVERING DNA TO MUSCLE CELLS  
USING RECOMBINANT ADENO-ASSOCIATED VIRUS VARIANTS

Description

10 Technical Field

The present invention relates generally to DNA delivery methods. More particularly, the invention relates to the use of recombinant adeno-associated virus (AAV) virions for delivery of a selected gene to muscle cells and tissue. The method provides for sustained, high-level expression of the delivered gene.

20 Background of the Invention

Gene delivery is a promising method for the treatment of acquired and inherited diseases. Muscle tissue is an appealing gene delivery target because it is readily accessible, well-differentiated and nondividing. Barr and Leiden (1991) *Science* 254:1507-1509. These properties are important in the selection of appropriate delivery strategies to achieve maximal gene transfer.

30 Several experimenters have demonstrated the ability to deliver genes to muscle cells with the subsequent systemic circulation of proteins encoded by the delivered genes. See, e.g., Wolff et al. (1990) *Science* 247:1465-1468; Accardi et al. (1991) *Nature* 352:815-818; Barr and Leiden (1991) *Science* 254:1507-1509; Dhanan et al. (1991) *Science* 254:1509-1512; Wolff et al. (1992) *Human Mol. Genet.* 1:361-369; Eyal et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:4523-4527; Davis et al. (1993) *Hum. Gene Therapy* 4:151-159.

-1-

5 Genes have been delivered to muscle by direct injection of plasmid DNA, such as described by Wolff et al. (1990) *Science* 247:1465-1468; Accardi et al. (1991) *Nature* 352:815-818; Barr and Leiden (1991) *Science* 254:1507-1509. However, this mode of administration generally results in sustained but low levels of expression. Low but sustained expression levels may be effective in certain situations, such as for providing immunity.

10 Viral based systems have also been used for gene delivery to muscle. For example, human adenoviruses are double-stranded DNA viruses which enter cells by receptor-mediated endocytosis. These viruses have been considered well suited for gene transfer because they are easy to grow and manipulate and they exhibit a broad host range *in vivo* and *in vitro*. Adenoviruses are able to infect quiescent as well as replicating target cells and persist extrachromosomally, rather than integrating into the host genome.

20 Despite these advantages, adenovirus vectors suffer from several drawbacks which make them ineffective for long term gene therapy. In particular, adenovirus vectors express viral proteins that may elicit an immune response which may decrease the life of the transduced cell. This immune reaction may preclude subsequent treatments because of humoral and/or T cell responses. Furthermore, the adult muscle cell may lack the receptor which recognizes adenovirus vectors, precluding efficient transduction of this cell type using such vectors.

30 Thus, attempts to use adenoviral vectors for the delivery of genes to muscle cells has resulted in poor and/or transitory expression. See, e.g., Quantin et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:2581-2584; Accardi et al. (1994) *Hum. Mol. Genetics* 3:579-584; Accardi et al. (1994) *Gene Therapy* 1:338-340; Dai et

-2-

a1. (1995) *Proc. Natl. Acad. Sci. USA* 92:1401-1405;  
Descamps et al. (1995) *Gene Therapy* 2:411-417;  
Clijnenkrantz et al. (1995) *Hum. Gene Therapy* 6:1265-  
1270.

5 Gene therapy methods based upon surgical  
transplantation of myoblasts has also been attempted.  
See, e.g., International Publication no. WO 95/13376;  
Dhavan et al. (1991) *Science* 253:1509-1512; Wolff et  
al. (1992) *Hum. Mol. Genet.* 1:363-369; Dai et al.  
10 (1992) *Proc. Natl. Acad. Sci. USA* 89:10892-10895;  
Kamamoto et al. (1994) *Hum. Gene Therapy* 5:1349-1356;  
Ramao et al. (1995) *J. Clin. Invest.* 95:1808-1813;  
Blau and Springer (1995) *New Eng. J. Med.* 333:1204-  
1207; Leiden, J.M. (1995) *New Eng. J. Med.* 333:871-  
15 872; Mendell et al. (1995) *New Eng. J. Med.* 333:832-  
838; and Blau and Springer (1995) *New Eng. J. Med.*  
333:1554-1556. However, such methods require  
substantial tissue culture manipulation and surgical  
expertise, and, at best, show inconclusive efficacy in  
20 clinical trials. Thus, a simple and effective method  
of gene delivery to muscle, resulting in long-term  
expression of the delivered gene, would be desirable.

Recombinant vectors derived from an adeno-  
associated virus (AAV) have been used for gene  
25 delivery. AAV is a helper-dependent DNA parvovirus  
which belongs to the genus Dependovirus. AAV requires  
infection with an unrelated helper virus, such as  
adenovirus, a herpesvirus or vaccinia, in order for a  
productive infection to occur. The helper virus  
30 supplies accessory functions that are necessary for  
most steps in AAV replication. In the absence of such  
infection, AAV establishes a latent state by insertion  
of its genome into a host cell chromosome. Subsequent  
infection by a helper virus rescues the integrated  
copy which can then replicate to produce infectious  
35 viral progeny. AAV has a wide host range and is able  
to replicate in cells from any species so long as

-3-

5 There is also a successful infection of such cells  
with a suitable helper virus. Thus, for example,  
human AAV will replicate in canine cells coinfectected  
with a canine adenovirus. AAV has not been associated  
with any human or animal disease and does not appear  
to alter the biological properties of the host cell  
upon integration. For a review of AAV, see, e.g.,  
Berns and Boehrsky (1987) *Advances in Virus Research*  
10 (Academic Press, Inc.) 32:243-307.

The AAV genome is composed of a linear,  
single-stranded DNA molecule which contains  
approximately 4681 bases (Berns and Boehrsky, supra).  
The genome includes inverted terminal repeats (ITRs)  
at each end which function in cis as origins of DNA  
15 replication and as packaging signals for the virus.  
The internal nonrepeated portion of the genome  
includes two large open reading frames, known as the  
AAV rep and cap regions, respectively. These regions  
code for the viral proteins involved in replication  
and packaging of the virion. For a detailed  
20 description of the AAV genome, see, e.g., Muzyczka, N.  
(1992) *Current Topics in Microbiol. and Immunol.*  
158:97-129.

The construction of recombinant AAV (rAAV)  
25 virions has been described. See, e.g., U.S. Patent  
Nos. 5,173,414 and 5,339,941; International  
Publication Numbers WO 92/01070 (published 23 January  
1992) and WO 93/03769 (published 4 March 1993);  
Lebkowski et al. (1988) *Molec. Cell. Biol.* 8:3588-  
30 3596; Vincent et al. (1990) *Vaccines* 96 (Cold Spring  
Harbor Laboratory Press); Carter, B.J. (1992) *Current  
Opinion in Biotechnology* 3:533-539; Muzyczka, N.  
(1992) *Current Topics in Microbiol. and Immunol.*  
158:97-129; and Kotlin, R.W. (1994) *Human Gene Therapy*  
35 5:793-801.

Recombinant AAV virion production generally  
involves cotransfection of a producer cell with an AAV

-4-

vector plasmid and a helper construct which provides AAV helper functions to complement functions missing from the AAV vector plasmid. In this manner, the producer cell is capable of expressing the AAV proteins necessary for AAV replication and packaging. The AAV vector plasmid will include the DNA of interest flanked by AAV ITRs which provide for AAV replication and packaging functions. AAV helper functions can be provided via an AAV helper plasmid that includes the AAV rep and/or cap coding regions but which lacks the AAV ITRs. Accordingly, the helper plasmid can neither replicate nor package itself. The producer cell is then infected with a helper virus to provide accessory functions, or with a vector which includes the necessary accessory functions. The helper virus transactivates the AAV promoters present on the helper plasmid that direct the transcription and translation of AAV rep and cap regions. Upon subsequent culture of the producer cells, recombinant AAV virions harboring the DNA of interest, are produced.

Recombinant AAV virions have been shown to exhibit tropism for respiratory epithelial cells (Plotte et al. (1992) *Am. J. Respir. Cell Mol. Biol.* 7:149-156; Plotte et al. (1993) *J. Biol. Chem.* 268:3781-3790; Plotte et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:10613-10617) and neurons of the central nervous system (Kapilite et al. (1994) *Nature Genetics* 8:148-154). These cell types are well-differentiated slowly-dividing or postmitotic. Plotte et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:10613-10617; Kapilite et al. (1994) *Nature Genetics* 8:148-154. The ability of AAV vectors to transduce nonproliferating cells (Podanskoff et al. (1994) *J. Virol.* 68:5656-5666; Russell et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:8915-8919; Plotte et al. (1994) *Am. J. Respir. Cell Mol. Biol.* 11:517-521) along with the attributes of

being inherently defective and nonpathogenic, place  $\Delta$ AV in a unique position among gene therapy viral vectors.

Despite these advantages, the use of recombinant AAV virions to deliver genes to muscle cells *in vivo* has not heretofore been disclosed.

### Summary of the Invention:

10 accordingly, the present invention is based  
on the surprising and unexpected discovery that  
recombinant AAV (rAAV) virions provide for efficient  
delivery of genes and sustained production of  
therapeutic proteins in various muscle cell types.  
The invention allows for *in vivo* secretion of the  
therapeutic protein from transduced muscle cells such  
that systemic delivery of therapeutic levels of the  
protein is achieved. These results are seen with both  
*in vivo* and *in vitro* modes of DNA delivery. Hence,  
rAAV virions allow delivery of DNA directly to muscle  
tissue. The ability to deliver and express genes in  
muscle cells, as well as to provide for secretion of  
the produced protein from transduced cells, allows the  
use of gene therapy approaches to treat and/or prevent  
a wide variety of disorders.

Furthermore, the ability to deliver DNA muscle cells by intramuscular administration *in vivo* provides a more efficient and convenient method of gene transfer.

Thus, in one embodiment, the invention relates to a method of delivering a selected gene to a muscle cell or tissue. The method comprises:

(a) providing a recombinant AAV vector which comprises an AAV vector comprising the selected gene operably linked to control elements capable of directing the *in vivo* transcription and translation of the selected gene; and

(b) Introducing the recombinant AAV virion into the muscle cell or tissue.

In particularly preferred embodiments, the selected gene encodes a therapeutic protein, such as erythropoietin (EPO), or the lysosomal enzyme, acid  $\alpha$ -glucosidase (GAA).

In another embodiment, the invention is directed to a muscle cell or tissue transduced with a recombinant AAV virion which comprises an AAV vector, the AAV vector comprising a selected gene operably linked to control elements capable of directing the *in vivo* transcription and translation of the selected gene.

In still further embodiments, the invention is directed to a method of treating an acquired or inherited disease in a mammalian subject comprising introducing into a muscle cell or tissue of the subject, *in vivo*, a therapeutically effective amount of a pharmaceutical composition which comprises (a) a recombinant AAV virions. The recombinant AAV virions comprise an AAV vector, the AAV vector comprising a selected gene operably linked to control elements capable of directing the transcription and translation of the selected gene when present in the subject.

In yet another embodiment, the invention is directed to a method of treating an acquired or inherited disease in a mammalian subject comprising:

(a) Introducing a recombinant AAV virion into a muscle cell or tissue *in vitro* to produce a transduced muscle cell. The recombinant AAV virion comprises an AAV vector, the AAV vector comprising a selected gene operably linked to control elements capable of directing the transcription and translation of the selected gene when present in the subject; and  
(b) administering to the subject a therapeutically effective amount of a composition

comprising a pharmaceutically acceptable excipient and the transduced muscle cells from step (a).

In a further embodiment, the invention relates to a method for delivering a therapeutically effective amount of a protein systemically to a mammalian subject comprising introducing into a muscle cell or tissue of the subject a pharmaceutical composition which comprises (a) a pharmaceutically acceptable excipient; and (b) recombinant AAV virions, wherein the recombinant AAV virions comprise an AAV vector, the AAV vector comprising a selected gene operably linked to control elements capable of directing the transcription and translation of the selected gene when present in the subject, wherein the introducing is done *in vivo*.

In another embodiment, the invention is directed to a method for delivering a therapeutically effective amount of a protein systemically to a mammalian subject comprising:

(a) Introducing a recombinant AAV virion into a muscle cell or tissue *in vitro* to produce a transduced muscle cell, wherein the recombinant AAV virion comprises an AAV vector, the AAV vector comprising a selected gene operably linked to control elements capable of directing the transcription and translation of the selected gene when present in the subject; and

(b) administering to the subject a therapeutically effective amount of a composition comprising a pharmaceutically acceptable excipient and the transduced muscle cells from step (a).

In other embodiments, the invention is directed to an AAV vector comprising a gene encoding either erythropoietin (EPO), or acid  $\alpha$ -glucosidase (GAA), operably linked to control elements capable of directing the *in vivo* transcription and translation of

the gene, as well as a recombinant AAV (rAAV) vector comprising the vector.

These and other embodiments of the subject invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

#### Brief Description of the Figures

Figure 1 shows *in situ* histochemical detection of  $\beta$ -galactosidase expression in murine muscle cells following transduction with rAAV-LacZ as described in Example 3, Part A. In the study, the tibialis anterior muscle of adult Balb/c mice was injected with  $8 \times 10^5$  rAAV-LacZ. Animals were sacrificed (a) 2, (b) 4, (c) 8, (d) 12, (e) 24, or (f) 32 weeks after injection. The tibialis anterior was excised, and 10  $\mu$ m sections were stained by X-gal for  $\beta$ -galactosidase histochemistry. The stained tissue samples were photographed at 25X.

Figure 2 shows a section of skeletal muscle two months after injection with rAAV-LacZ as described in Example 3, Part A. The tibialis anterior muscle was processed for *in situ* detection of  $\beta$ -galactosidase expression, and photographed with diffraction-interference contrast optics at 400X.

Figure 3 depicts  $\beta$ -galactosidase expression in Balb/c mice tibialis anterior muscle transduced *in vivo* with rAAV-LacZ as described in Example 3, Part B. Adult Balb/c mice were injected intramuscularly (IM) with various doses of rAAV-LacZ. At 2 and 8 weeks post injection, tissue was harvested for analysis of  $\beta$ -galactosidase ( $\beta$ -gal).  $\beta$ -gal expression was analyzed by measurement of relative light units (RLU) emitted from muscle homogenates, as detected by luminometer.

Figure 4 shows the secretion of human erythropoietin (hEPO) from transduced myotubes and myoblasts, as described in Example 4. Myotubes

(differentiated cells) or myoblasts (actively dividing cells) were transduced with rAAV-hEPO at a ratio of approximately  $10^5$  per target cell. Levels of secreted hEPO were analyzed in supernatants at various time points. Baseline levels of hEPO prior to transduction were below the level of detection in both cell populations; the values at each time point represent replicate values  $\pm$  standard deviation.

Figure 5 shows the secretion of human erythropoietin (hEPO) by C2C12 myotubes transduced with rAAV-hEPO as described in Example 4. Confluent C2C12 myoblasts were differentiated into myotubes and transduced with  $3 \times 10^8$  (open bar),  $3 \times 10^9$  (cross-hatched bar), or  $3 \times 10^{10}$  (solid bar) rAAV-hEPO. Secretion of EPO was measured 3, 8, and 14 days after transduction. Control rAAV-LacZ myotubes secreted  $<2.5$  mU/mL EPO. The bar graph depicts mean production of EPO/well/24 hour as determined in triplicate cultures  $\pm$  the standard error of mean (SEM).

Figure 6 shows the secretion of human erythropoietin (hEPO) by primary human myotubes transduced with rAAV-hEPO as described in Example 5. Confluent human myoblasts were differentiated into myotubes by culture for 14 days in reduced-serum media, then transduced with  $3 \times 10^8$  (open bar),  $3 \times 10^9$  (cross-hatched bar), or  $3 \times 10^{10}$  (solid bar) rAAV-hEPO. Secretion of EPO was measured 3, 8 and 14 days after transduction. Control myotubes transduced with rAAV-LacZ secreted  $<2.5$  mU/mL EPO. The bar graph depicts mean production of EPO/well/24 hour as determined in triplicate cultures  $\pm$  SEM.

Figure 7 depicts the time course of EPO secretion in Balb/c mice after IM injection with rAAV-hEPO. Adult Balb/c mice were injected IM with  $1 \times 10^{10}$  (V),  $3 \times 10^{10}$  (A),  $1 \times 10^{11}$  (B), or  $3 \times 10^{11}$  (C) purified rAAV-LacZ at day = 0, and serum EPO levels

measured at various time points after injection. Reported values represent means ( $n=4$ )  $\pm$  SEM.

Figure 8 shows high level expression of acid  $\alpha$ -glucosidase (GAA) in human skeletal muscle transduced *in vitro* with rAAV-hGAA as described in Example 8, Part A. In the study, differentiated human myoblasts were exposed to rAAV-hGAA virions at a MOI of  $2 \times 10^5$ . Cells were collected at the time points indicated, and GAA activity measured by enzymatic assay. Non-transduced control cells (open bar) and cells transduced with rAAV-LacZ (cross-hatched bar) showed no significant expression of GAA, while cells transduced with rAAV-hGAA (solid bar) showed high levels of GAA activity. The bar graph represents mean GAA activity determined in triplicate cultures  $\pm$  SEM. Figure 9 shows expression of acid  $\alpha$ -glucosidase in Balb/c mice tibialis anterior muscle cells that were transduced *in vivo* with rAAV-hGAA, as described in Example 8, Part B. Adult Balb/c mice were injected intramuscularly (IM) with  $4 \times 10^{13}$  rAAV-hGAA (solid bar) or the same dose of rAAV-LacZ (open bar). At various time points after injection, muscle tissue was harvested for analysis of GAA activity by enzymatic assay. The bar graph shows mean GAA activity determined in five animals (weeks 1 and 4), or in four animals (week 10)  $\pm$  SEM.

#### Detailed Description of the Invention

The practice of the present invention will employ, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, et al. *Molecular Cloning: A Laboratory Manual* (Current Edition); DNA Cloning: A Practical Approach, vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed.,

Current Edition); *Nucleic Acid Hybridization* (B. Hanes & S. Higgins, eds., Current Edition); *Transcription and Translation* (B. Hanes & S. Higgins, eds., Current Edition); *CRC Handbook of Parvoviruses*, vol. I & II (P. Tijssen, ed.); *Fundamental Virology*, 2nd Edition, vol. I & II (B.K. Fields and D.R. Knipe, eds.).

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content clearly dictates otherwise.

#### A. Definitions

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

The phrase "gene delivery" or "gene transfer" refers to methods or systems for reliably inserting foreign DNA into target cells, such as into muscle cells. Such methods can result in transient or long term expression of genes. Gene transfer provides a unique approach for the treatment of acquired and inherited diseases. A number of systems have been developed for gene transfer into mammalian cells. See, e.g., U.S. Patent No. 5,399,346.

The term "therapeutic protein" refers to a protein which is defective or missing from the subject in question, thus resulting in a disease state or disorder in the subject, or to a protein which confers a benefit to the subject in question, such as an antiviral, antibacterial or antitumor function. A therapeutic protein can also be one which modifies any one of a wide variety of biological functions, such as endocrine, immunological and metabolic functions. Representative therapeutic proteins are discussed more fully below.

By "vector" is meant any genetic element, such as a plasmid, phage, transposon, cosmid,

chromosome, virus, virion, etc., which is capable of replication when associated with the proper control elements and which can transfer gene sequences between cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

By "AAV vector" is meant a vector derived from an adeno-associated virus serotype, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAVX7, etc. AAV vectors can have one or more of the AAV wild-type genes deleted in whole or part, preferably the *rep* and/or *cap* genes (described below), but retain functional flanking ITR sequences (also described below). Functional ITR sequences are necessary for the rescue, replication and packaging of the AAV virion. Thus, an AAV vector is defined herein to include at least those sequences required in *cis* for replication and packaging (e.g., functional ITRs) of the virus. The ITRs need not be the wild-type nucleotide sequences, and may be altered, e.g., by the insertion, deletion or substitution of nucleotides, so long as the sequences provide for functional rescue, replication and packaging.

By "recombinant virus" is meant a virus that has been genetically altered, e.g., by the addition or insertion of a heterologous nucleic acid construct into the particle.

By "AAV virion" is meant a wild-type (wt) AAV virus particle (comprising a linear, single-stranded AAV nucleic acid genome associated with an AAV capsid protein coat). In this regard, single-stranded AAV nucleic acid molecules of either complementary sense, e.g., "sense" or "antisense" strands, can be packaged into any one AAV virion and both strands are equally infectious.

A "recombinant AAV virion," or "rAAV virion" is defined herein as an infectious, replication-defective virus composed of an AAV protein shell,

encapsidating a DNA molecule of interest which is flanked on both sides by AAV ITRs. An rAAV virion is produced in a suitable producer cell which has had an AAV vector, AAV helper functions and accessory functions introduced therein. In this manner, the producer cell is rendered capable of encoding AAV polypeptides that are required for packaging the AAV vector (containing a recombinant nucleotide sequence of interest) into recombinant virion particles for subsequent gene delivery.

The term "transfection" is used to refer to the uptake of foreign DNA by a mammalian cell. A cell has been "transfected" when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are known in the art. See, e.g., Graham et al. (1973) *Virology*, 52:456, Sambrook et al. (1989) *Molecular Cloning, a laboratory manual*, Cold Spring Harbor Laboratory, New York, Davis et al. (1986) *Basic Methods in Molecular Biology*, Elsevier, and Chu et al. (1983) *Gene* 13:197. Such techniques can be used to introduce one or more exogenous DNA moieties, such as a plasmid vector and other nucleic acid molecules, into suitable cells. The term refers to both stable and transient uptake of the genetic material.

The term "transduction" denotes the delivery of a DNA molecule to a recipient cell either *in vivo* or *in vitro*, via a replication-defective viral vector, such as via a recombinant AAV virion.

By "muscle cell" or "tissue" is meant a cell or group of cells derived from muscle, including but not limited to cells and tissue derived from skeletal muscle; smooth muscle, e.g., from the digestive tract, urinary bladder and blood vessels; and cardiac muscle. The term captures muscle cells both *in vitro* and *in vivo*. Thus, for example, an isolated cardiomyocyte would constitute a "muscle cell" for purposes of the



present invention, as would a muscle cell as it exists in muscle tissue present in a subject *in vivo*. The term also encompasses both differentiated and nondifferentiated muscle cells, such as myocytes, myotubes, myoblasts, cardiomyocytes and cardiomyoblasts.

The term "heterologous" as it relates to nucleic acid sequences such as gene sequences and control sequences, denotes sequences that are not normally joined together, and/or are not normally associated with a particular cell. Thus, a "heterologous" region of a nucleic acid construct or a vector is a segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule *in nature*. For example, a heterologous region of a nucleic acid construct could include a coding sequence flanked by sequences not found in association with the coding sequence *in nature*. Another example of a heterologous coding sequence is a construct in which the coding sequence itself is not found *in nature* (e.g., synthetic sequences having codons different from the native gene). Similarly, a cell transformed with a construct which is not normally present in the cell would be considered heterologous for purposes of this invention. Allelic variation or naturally occurring mutational events do not give rise to heterologous DNA, as used herein.

By "DNA" is meant a polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in double-stranded or single-stranded form, either relaxed and supercoiled. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes single- and double-stranded DNA found, *inter alia*, in linear DNA molecules (e.g., restriction fragments), viruses,

plasmids, and chromosomes. In discussing the structure of particular DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (i.e., the strand having the sequence homologous to the mRNA). The term captures molecules that include the four bases adenine, guanine, thymine, or cytosine, as well as molecules that include base analogues which are known in the art.

A "gene" or "coding sequence" or a sequence which "encodes" a particular protein, is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the gene are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A gene can include, but is not limited to, cDNA from prokaryotic or eukaryotic mRNA, genomic DNA sequences from prokaryotic or eukaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence will usually be located 3' to the gene sequence.

The term "control elements" refers collectively to promoter regions, polyadenylation signals, transcription termination sequences, upstream regulatory domains, origins of replication, internal ribosome entry sites ("IRES"), enhancers, and the like, which collectively provide for the replication, transcription and translation of a coding sequence in a recipient cell. Not all of these control elements need always be present so long as the selected coding sequence is capable of being replicated, transcribed and translated in an appropriate host cell.

The term "promoter region" is used herein in its ordinary sense to refer to a nucleotide region

comprising a DNA regulatory sequence, wherein the regulatory sequence is derived from a gene which is capable of binding RNA polymerase and initiating transcription of a downstream (3'-direction) coding sequence.

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control elements operably linked to a coding sequence are capable of effecting the expression of the coding sequence. The control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

For the purpose of describing the relative position of nucleotide sequences in a particular nucleic acid molecule throughout the instant application, such as when a particular nucleotide sequence is described as being situated "upstream," "downstream," "3'," or "5'" relative to another sequence, it is to be understood that it is the position of the sequences in the "sense" or "coding" strand of a DNA molecule that is being referred to as is conventional in the art.

"Homology" refers to the percent of identity between two polynucleotide or two polypeptide moieties. The correspondence between the sequence from one moiety to another can be determined by techniques known in the art. For example, homology can be determined by a direct comparison of the sequence information between two polypeptide molecules by aligning the sequence information and using readily available computer programs. Alternatively, homology

can be determined by hybridization of polynucleotides under conditions which form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s), and size determination of the digested fragments. Two DNA, or two polypeptide sequences are "substantially homologous" to each other when at least about 80%, preferably at least about 90%, and most preferably at least about 95% of the nucleotides or amino acids match over a defined length of the molecules, as determined using the methods above.

By "mammalian subject" is meant any member of the class Mammalia including, without limitation, humans and nonhuman primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs, and the like. The term does not denote a particular age or sex. Thus, adult and newborn subjects, as well as fetuses, whether male or female, are intended to be covered.

#### B. General Methods

The present invention provides for the successful transfer of a selected gene to a muscle cell using recombinant AAV virions. The method allows for the direct, *in vivo* injection of recombinant AAV virions into muscle tissue, e.g., by intramuscular injection, as well as for the *in vitro* transduction of muscle cells which can subsequently be introduced into a subject for treatment. The invention also provides for secretion of the produced protein *in vivo*, from transduced muscle cells, such that systemic delivery can be achieved.

Muscle provides a desirable target for gene therapy since muscle cells are readily accessible and

nondividing. However, the present invention also finds use with nondifferentiated muscle cells, such as myoblasts, which can be transduced *in vitro*, and subsequently introduced into a subject.

Since muscle has ready access to the circulatory system, a protein produced and secreted by muscle cells and tissue *in vivo* will enter the bloodstream for systemic delivery. Furthermore, since sustained, therapeutic levels of protein secretion from muscle is achieved *in vivo* using the present invention, repeated parenteral delivery is avoided or reduced in frequency such that therapy can be accomplished using only one or few injections. Thus, the present invention provides significant advantages over prior gene delivery methods.

The recombinant AAV virions of the present invention, including the DNA of interest, can be produced using standard methodology, known to those of skill in the art. The methods generally involve the steps of (1) introducing an AAV expression vector into a producer cell; (2) introducing an AAV helper construct into the producer cell, where the helper construct includes AAV coding regions capable of being expressed in the producer cell to complement AAV helper functions missing from the AAV vector; (3) introducing one or more helper viruses and/or accessory function vectors into the producer cell, wherein the helper virus and/or accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the cell; and (4) culturing the producer cell to produce rAAV virions. The AAV expression vector, AAV helper construct and the helper virus or accessory function vector(s) can be introduced into the producer cell, either simultaneously or serially, using standard transfection techniques.

#### 1. AAV Expression Vectors

AAV expression vectors are constructed using known techniques to at least provide as operatively linked components in the direction of transcription, control elements including a transcriptional initiation region, the DNA of interest and a transcriptional termination region. The control elements are selected to be functional in a mammalian muscle cell. The resulting construct which contains the operatively linked components is bounded (5' and 3') with functional AAV ITR sequences.

The nucleotide sequences of AAV ITR regions are known. See, e.g., Kotlik, R.N. (1994) *Human Gene Therapy* 5:753-801; Berns, K.I. "Parvoviridae and their replication" in *Fundamental Virology*, 2nd Edition, (B.N. Fields and D.M. Knipe, eds.) for the AAV-2 sequence. AAV ITRs used in the vectors of the invention need not have a wild-type nucleotide sequence, and may be altered, e.g., by the insertion, deletion or substitution of nucleotides.

Additionally, AAV ITRs may be derived from any of several AAV serotypes, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAVX, etc. Furthermore, 5' and 3' ITRs which flank a selected nucleotide sequence in an AAV expression vector need not necessarily be identical or derived from the same AAV serotype or isolate, so long as they function as intended, i.e., to allow for packaging of virions.

Suitable DNA molecules for use in AAV vectors will be less than about 5 kilobases (kb) in size and will include, for example, a gene that encodes a protein that is defective or missing from a recipient subject or a gene that encodes a protein having a desired biological or therapeutic effect (e.g., an antibacterial, antiviral or antitumor function).

Suitable DNA molecules include, but are not limited to, those encoding for proteins used for the treatment of endocrine, metabolic, hematologic, cardiovascular, neurologic, musculoskeletal, urologic, pulmonary and immune disorders, including such disorders as inflammatory diseases, autoimmune, chronic and infectious diseases, such as AIDS, cancer, hypercholesterolemia, insulin disorders such as diabetes, growth disorders, various blood disorders including various anemias, thalassemias and hemophilia; genetic defects such as cystic fibrosis, Gaucher's Disease, Hunter's Disease, adenosine deaminase (ADA) deficiency, epiphysema, or the like.

To exemplify the invention, the gene encoding erythropoietin (EPO) can be used. EPO is a glycoprotein hormone produced in fetal liver and adult kidney which acts on progenitor cells in the bone marrow and other hematopoietic tissue to stimulate the formation of red blood cells. Genes encoding human and other mammalian EPO have been cloned, sequenced and expressed, and show a high degree of sequence homology in the coding region across species. Wen et al. (1993) *Glucol* 82:1507-1516. The sequence of the gene encoding native human EPO, as well as methods of obtaining the same, are described in, e.g., U.S. Patent Nos. 4,954,437 and 4,703,008, as well as in Jacobs et al. (1985) *Nature* 313:806-810; Lin et al. (1985) *Proc. Natl. Acad. Sci. USA* 82:7580; International Publication Number WO 85/03610; and European Patent Publication Number 232,034 B2. In addition, the sequences of the genes encoding native feline, canine and porcine EPO are known and readily available (GenBank Accession Nos.: L10606; L13027; and L10607, respectively), and the sequence of the gene encoding monkey (*Macaca mulatta*) is also known and available (GenBank Accession No.: L10609). The term "EPO" as used herein refers to the native, full-length

-21-

secreted form of EPO, as well as to analogs or derivatives thereof comprising single or multiple amino acid substitutions, deletions or additions which retain EPO function or activity. In this regard, a number of small peptides have been identified which bind to and activate the receptor for EPO. Wrighton et al. (1995) *Science* 223:456-463; Livanh et al. (1996) *Science* 223:464-471. The recombinant MAV virions described herein which include a gene encoding EPO, or encoding an analog or derivative thereof having the same function, are particularly useful in the treatment of blood disorders characterized by defective red blood cell formation, such as in the treatment of anemia. Increased red blood cell production due to the production of EPO can be readily determined by an appropriate indicator, such as by comparing hematocrit measurements pre- and post-treatment, measuring increases in red blood cell count, hemoglobin concentration, or in reticulocyte counts. As described above, the EPO gene is flanked by MAV ITRs.

Alternatively, a nucleotide sequence encoding the lysosomal enzyme acid  $\alpha$ -glucosidase (GAA) can be used. GAA functions to cleave  $\alpha$ -1,4 and  $\alpha$ -1,6 linkages of lysosomal glycogen to release monosaccharides. The sequence of the gene encoding human GAA, as well as methods of obtaining the same, have been previously described (GenBank Accession Numbers: M34424 and Y00639; Martiniuk et al. (1990) *DNA Cell Biol.* 2:85-94; Martiniuk et al. (1986) *Proc. Natl. Acad. Sci. USA* 83:9641-9644; Neufeld et al. (1988) *Eur. Mol. Biol. Organ.* 2:1697-1704). Thus, the recombinant MAV virions described herein can include a nucleotide sequence encoding GAA, or encoding an analog or derivative thereof having GAA activity.

The selected nucleotide sequence, such as EPO or another gene of interest, is operably linked to

-22-

control elements that direct the transcription or expression thereof in the subject *in vivo*. Such control elements can comprise control sequences normally associated with the selected gene.

Alternatively, heterologous control sequences can be employed. Useful heterologous control sequences generally include those derived from sequences encoding mammalian or viral genes. Examples include, but are not limited to, the SV40 early promoter; mouse mammary tumor virus LTR promoter; adenovirus major late promoter (Ad MLP); herpes simplex virus (HSV) promoters; a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter region (CMVIE); a Rous sarcoma virus (RSV) promoter; synthetic promoters; hybrid promoters; and the like. In addition, sequences derived from nonviral genes, such as the murine metallothionein gene, will also find use herein. Such promoter sequences are commercially available from, e.g., Stratagene (San Diego, CA).

For purposes of the present invention, control elements, such as muscle-specific and inducible promoters, enhancers and the like, will be of particular use. Such control elements include, but are not limited to, those derived from the actin and myosin gene families, such as from the myosin family (Kelntraub et al. (1991) *Science* 251:761-766); the myocyte-specific enhancer binding factor MEF-2 (Cserjesi and Olson (1991) *Mol. Cell Biol.* 11:4854-4862); control elements derived from the human skeletal actin gene (Nucast et al. (1987) *Mol. Cell Biol.* 7:4089-4099) and the cardiac actin gene; muscle creatine kinase sequence elements (Johnson et al. (1989) *Mol. Cell Biol.* 9:3393-3399) and the murine creatine kinase enhancer (MCK) element; control elements derived from the skeletal fast-twitch tropomyosin C gene, the slow-twitch cardiac tropomyosin C gene and the slow-twitch tropomyosin I gene; hypoxia-

inducible nuclear factors (Semenza et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:5680-5684; Semenza et al. *J. Biol. Chem.* 269:23737-23763); steroid-inducible elements and promoters, such as the glucocorticoid response element (GRE) (Kader and White (1993) *Proc. Natl. Acad. Sci. USA* 90:5607-5607); the fusion consensus element for RU486 induction; elements that provide for tetracycline regulated gene expression (Dhawan et al. (1995) *Somat. Cell. Mol. Genet.* 21:233-240; Shockett et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6522-6526; and inducible, synthetic humanized promoters (Rivera et al. (1996) *Nature Med.* 2:1028-1032).

These and other regulatory elements can be tested for potential *in vivo* efficacy using the *in vitro* myoblast model, which mimics quiescent *in vivo* muscle physiology, described in the examples below.

The AAV expression vector which harbors the DNA molecule of interest bounded by AAV ITRs, can be constructed by directly inserting the selected sequence(s) into an AAV genome which has had the major AAV open reading frames ("ORFs") excised therefrom. Other portions of the AAV genome can also be deleted, so long as a sufficient portion of the ITRs remain to allow for replication and packaging functions. Such constructs can be designed using techniques well known in the art. See, e.g., U.S. Patent Nos. 5,173,414 and 5,139,941; International Publication Nos. WO 92/02070 (published 23 January 1992) and WO 93/03769 (published 4 March 1993); Lebowicki et al. (1988) *Molec. Cell Biol.* 8:3988-3996; Vincent et al. (1993) *Vaccines* 90 (Cold Spring Harbor Laboratory Press); Carter, B.J. (1992) *Current Opinion in Biotechnology* 1:533-539; Nuzyczka, K. (1992) *Current Topics in Microbiol. and Immunol.* 158:97-129; Kotlin, R.M. (1994) *Human Gene Therapy* 5:793-801; Sheelling and Smith (1994) *Gene*

Therapy 1:165-169; and Zhou et al. (1994) *J. Exp. Med.* 179:1867-1875.

Alternatively, AAV ITRs can be excised from the viral genome or from an AAV vector containing the same and fused 5' and 3' of a selected nucleic acid construct that is present in another vector using standard ligation techniques, such as those described in Sambrook et al., supra. For example, ligations can be accomplished in 20 mM Tris-Cl pH 7.5, 10 mM MgCl<sub>2</sub>, 10 mM DTT, 33 µg/ml BSA, 10 mM-50 mM NaCl, and either 40 µM ATP, 0.01-0.02 (Waters) units T4 DNA ligase at 0°C (for "sticky end" ligation) or 1 mM ATP, 0.3-0.6 (Waters) units T4 DNA ligase at 14°C (for "blunt end" ligation). Inter-molecular "sticky end" ligations are usually performed at 30-100 µg/ml total DNA concentrations (5-100 nM total end concentration).

AAV vectors which contain ITRs have been described in, e.g., U.S. Patent no. 5,139,941. In particular, several AAV vectors are described therein which are available from the American Type Culture Collection ("ATCC") under Accession Numbers 53222, 53223, 53224, 53225 and 53226.

Additionally, chimeric genes can be produced synthetically to include AAV ITR sequences arranged 5' and 3' of one or more selected nucleic acid sequences. Preferred codons for expression of the chimeric gene sequence in mammalian muscle cells can be used. The complete chimeric sequence is assembled from overlapping oligonucleotides prepared by standard methods. See, e.g., Edge, *Nature* (1981) 292:755; Nambair et al. *Science* (1984) 223:1299; Jay et al. *J. Biol. Chem.* (1984) 259:6311.

In order to produce rAAV virions, an AAV expression vector is introduced into a suitable producer cell using known techniques, such as by transfection. A number of transfection techniques are generally known in the art. See, e.g., Graham et al.

(1973) *Virology* 52:456, Sambrook et al. (1989) *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratories, New York, Davis et al. (1986) *Basic Methods in Molecular Biology*, Elsevier, and Chu et al. (1981) *Gene* 12:197. Particularly suitable transfection methods include calcium phosphate co-precipitation (Graham et al. (1973) *Virology* 52:456-467), direct micro-injection into cultured cells (Capechi, M.R. (1980) *Cell* 22:479-488), electroporation (Shigemura et al. (1988) *BioTechniques* 6:742-751), liposome mediated gene transfer (Mannino et al. (1988) *BioTechniques* 6:682-690), lipid-mediated transduction (Felgner et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:7413-7417), and nucleic acid delivery using high-velocity microprojectiles (Klein et al. (1987) *Nature* 322:70-73).

For the purposes of the invention, suitable producer cells for producing rAAV virions include microorganisms, yeast cells, insect cells, and mammalian cells, that can be, or have been, used as recipients of a heterologous DNA molecule. The term includes the progeny of the original cell which has been transfected. Thus, a "producer cell" as used herein generally refers to a cell which has been transfected with an exogenous DNA sequence. Cells from the stable human cell line, 293 (readily available through, e.g., the American Type Culture Collection under Accession Number ATCC CRL1573) are preferred in the practice of the present invention. Particularly, the human cell line 293 is a human embryonic kidney cell line that has been transformed with adenovirus type-5 DNA fragments (Graham et al. (1977) *J. Gen. Virol.* 36:591, and expresses the adenoviral E1a and E1b genes (Aiello et al. (1979) *Virology* 94:460). The 293 cell line is readily transfected, and provides a particularly convenient platform in which to produce rAAV virions.

## 2. AAV Helper Functions

Producer cells containing the above-described AAV expression vectors must be rendered capable of providing AAV helper functions in order to replicate and encapsidate the nucleotide sequences flanked by the AAV ITRs to produce rAAV virions. AAV helper functions are generally AAV-derived coding sequences which can be expressed to provide AAV gene products that, in turn, function in *trans* for productive AAV replication. AAV helper functions are used herein to complement necessary AAV functions that are missing from the AAV expression vectors. Thus, AAV helper functions include one, or both of the major AAV ORFs, namely the *rep* and *cap* coding regions, or functional homologues thereof.

By "AAV rep coding region" is meant the art-recognized region of the AAV genome which encodes the replication proteins Rep 78, Rep 68, Rep 52 and Rep 40. These Rep expression products have been shown to possess many functions, including recognition, binding and nicking of the AAV origin of DNA replication, DNA helicase activity and modulation of transcription from AAV (or other heterologous) promoters. The Rep expression products are collectively required for replicating the AAV genome. For a description of the AAV rep coding region, see, e.g., Kuzyszka, H. (1992) *Current Topics in Microbiol. and Immunol.* 358:97-125; and Kotlin, R.H. (1994) *Human Gene Therapy* 5:793-802. Suitable homologues of the AAV rep coding region include the human herpesvirus 6 (HHV-6) rep gene which is also known to mediate AAV-2 DNA replication (Thomson et al. (1994) *Virology* 204:304-311).

By "AAV cap coding region" is meant the art-recognized region of the AAV genome which encodes the capsid proteins VP1, VP2, and VP3, or functional homologues thereof. These cap expression products are the capsid proteins which are collectively required

-27-

for packaging the viral genome. For a description of the AAV cap coding region, see, e.g., Kuzyszka, H. and Kotlin, R.H. (*supra*).

AAV helper functions are introduced into the producer cell by transfecting the cell with an AAV helper construct either prior to, or concurrently with, the transfection of the AAV expression vector. AAV helper constructs are thus used to provide at least transient expression of AAV rep and/or cap genes to complement missing AAV functions that are necessary for productive AAV infection. AAV helper constructs lack AAV ITRs and can neither replicate nor package themselves.

These constructs can be in the form of a plasmid, phage, transposon, cosmid, virus, or virion. A number of AAV helper constructs have been described, such as the commonly used plasmids pAAV/Ad and pIM29+45 which encode both Rep and Cap expression products. See, e.g., Samiaski et al. (1989) *J. Virol.* 63:3822-3828; and McCarty et al. (1991) *J. Virol.* 65:2936-2945. A number of other vectors have been described which encode Rep and/or Cap expression products. See, e.g., U.S. Patent No. 5,139,941.

Both AAV expression vectors and AAV helper constructs can be constructed to contain one or more optional selectable markers. Suitable markers include genes which confer antibiotic resistance or sensitivity to, impart color to, or change the antigenic characteristics of those cells which have been transfected with a nucleic acid construct containing the selectable marker when the cells are grown in an appropriate selective medium. Several selectable marker genes that are useful in the practice of the invention include the hygromycin B resistance gene (encoding aminoglycoside phosphotransferase (APH)) that allows selection in mammalian cells by conferring resistance to G418

-28-

(available from Sigma, St. Louis, Mo.). Other suitable markers are known to those of skill in the art.

## 5 2. Accessory Functions

The producer cell must also be rendered capable of providing non AAV derived functions, or "accessory functions," in order to produce rAAV virions. Accessory functions are non AAV derived viral and/or cellular functions upon which AAV is dependent for its replication. Thus, accessory functions include at least those non AAV proteins and RNAs that are required in AAV replication, including those involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of cap expression products and AAV capsid assembly. Viral-based accessory functions can be derived from any of the known helper viruses.

20 Particularly, accessory functions can be introduced into and then expressed in producer cells using methods known to those of skill in the art. Commonly, accessory functions are provided by infection of the producer cells with an unrelated helper virus. A number of suitable helper viruses are known, including adenoviruses; herpesviruses such as herpes simplex virus types 1 and 2; and vaccinia viruses. Nonviral accessory functions will also find use herein, such as those provided by cell

30 synchronization using any of various known agents. See, e.g., Buller et al. (1981) *J. Virol.* 40:241-247; McPherson et al. (1985) *Virology* 147:217-222; Schlehofer et al. (1986) *Virology* 152:115-117.

Alternatively, accessory functions can be provided using an accessory function vector. Accessory function vectors include nucleotide sequences that provide one or more accessory

-29-

functions. An accessory function vector is capable of being introduced into a suitable producer cell in order to support efficient AAV virion production in the cell. Accessory function vectors can be in the form of a plasmid, phage, transposon or cosmid. Accessory vectors can also be in the form of one or more linearized DNA or RNA fragments which, when associated with the appropriate control elements and enzymes, can be transcribed or expressed in a producer cell to provide accessory functions.

Nucleic acid sequences providing the accessory functions can be obtained from natural sources, such as from the genome of an adenovirus particle, or constructed using recombinant or synthetic methods known in the art. In this regard, adenovirus-derived accessory functions have been widely studied, and a number of adenovirus genes involved in accessory functions have been identified and partially characterized. See, e.g., Carter, B.J. (1990) "Adeno-Associated Virus Helper Functions," in *CRC Handbook of Parvoviruses*, vol. 1 (P. Tijssen, ed.), and Muzyczka, N. (1992) *Curr. Topics Microbiol. and Immun.* 158:97-129. Specifically, early adenoviral gene regions E1a, E1b, E4, and E1a and, possibly, E1b are thought to participate in the accessory process. Janik et al. (1981) *Proc. Natl. Acad. Sci. USA* 78:1925-1929. Herpesvirus-derived accessory functions have been described. See, e.g., Young et al. (1979) *Prog. Med. Virol.* 25:113. Vaccinia virus-derived accessory functions have also been described. See, e.g., Carter, B.J. (1990) *supra*, Schlehofer et al. (1986) *Virology* 152:110-117.

As a consequence of the infection of the producer cell with a helper virus, or transfection of the producer cell with an accessory function vector, accessory functions are expressed which transactivate the AAV helper construct to produce AAV Rep and/or cap

-30-



proteins. The Rep expression products excise the recombinant DNA (including the DNA of interest) from the rAAV expression vector. The Rep proteins also serve to duplicate the AAV genome. The expressed Cap proteins assemble into capsids, and the recombinant AAV genome is packaged into the capsids. Thus, productive AAV replication ensues, and the DNA is packaged into rAAV virions.

Following recombinant AAV replication, rAAV virions can be purified from the producer cell using a variety of conventional purification methods, such as CsCl gradients. Further, if infection is employed to express the accessory functions, residual helper virus can be inactivated, using known methods. For example, adenovirus can be inactivated by heating to temperatures of approximately 60°C for, e.g., 20 minutes or more. This treatment effectively inactivates only the helper virus since AAV is extremely heat stable while the helper adenovirus is heat labile.

The resulting rAAV virions are then ready for use for DNA delivery, such as in gene therapy applications, for the production of transgenic animals, in vaccination, and particularly for the delivery of genes to a variety of muscle cell types.

#### 4. In vitro and in vivo delivery of rAAV virions

Generally, rAAV virions are introduced into a muscle cell using either *in vivo* or *in vitro* transduction techniques. If transduced *in vitro*, the desired recipient muscle cell will be removed from the subject, transduced with rAAV virions and reintroduced into the subject. Alternatively, syngeneic or xenogeneic muscle cells can be used where those cells will not generate an inappropriate immune response in the subject.

Suitable methods for the delivery and introduction of transduced cells into a subject have been described. For example, cells can be transduced *in vitro* by combining recombinant AAV virions with muscle cells e.g., in appropriate media, and screening for those cells harboring the DNA of interest using conventional techniques such as Southern blots and/or PCR, or by using selectable markers. Transduced cells can then be formulated into pharmaceutical compositions, described more fully below, and the composition introduced into the subject by various techniques, such as by intramuscular, intravenous, subcutaneous and intraperitoneal injection, or by injection into smooth and cardiac muscle, using e.g., a catheter.

For *in vivo* delivery, the rAAV virions will be formulated into pharmaceutical compositions and will generally be administered parenterally, e.g., by intramuscular injection directly into skeletal or cardiac muscle.

Pharmaceutical compositions will comprise sufficient genetic material to produce a therapeutically effective amount of the protein of interest, i.e., an amount sufficient to reduce or ameliorate symptoms of the disease state in question or an amount sufficient to confer the desired benefit. The pharmaceutical compositions will also contain a pharmaceutically acceptable excipient. Such excipients include any pharmaceutical agent that does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, glycerol and ethanol. Pharmaceutically acceptable salts can be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides,

phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. A thorough discussion of pharmaceutically acceptable excipients is available in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J., 1991).

Appropriate doses will depend on the mammal being treated (e.g., human or nonhuman primate or other mammal), age and general condition of the subject to be treated, the severity of the condition being treated, the particular therapeutic protein in question, its mode of administration, among other factors. An appropriate effective amount can be readily determined by one of skill in the art.

Thus, a "therapeutically effective amount" will fall in a relatively broad range that can be determined through clinical trials. For example, for *in vivo* injection, i.e., injection directly to skeletal or cardiac muscle, a therapeutically effective dose will be on the order of from about  $10^1$  to  $10^{15}$  of the rAAV virions, more preferably  $10^3$  to  $10^{14}$  rAAV virions. For *in vitro* transduction, an effective amount of rAAV virions to be delivered to muscle cells will be on the order of  $10^5$  to  $10^{12}$  of the rAAV virions. The amount of transduced cells in the pharmaceutical compositions will be from about  $10^4$  to  $10^{10}$  muscle cells, more preferably  $10^5$  to  $10^8$  muscle cells. When the transduced cells are introduced to vascular smooth muscle, a lower dose may be appropriate. Other effective dosages can be readily established by one of ordinary skill in the art through routine trials establishing dose response curves.

-33-

Dosage treatment may be a single dose schedule or a multiple dose schedule. Moreover, the subject may be administered as many doses as appropriate. One of skill in the art can readily determine an appropriate number of doses.

### C. Experimental

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

### Materials and Methods

#### 20 Vector constructs

##### A. Construction of p1903adhlacZ.

Plasmid p1903adhlacZ was used as the helper construct in the following examples and was constructed from plasmid pAdhlacZ. Plasmid pAdhlacZ was constructed by partially digesting plasmid pUC19 (GeneBank Reference Name: U07649, GeneBank Accession Number: U07649) with *Acl*III and *Bsp*HI, blunt-end modifying with the *Klenow* enzyme, and then ligating to form a circular 1732 bp plasmid containing the bacterial origin and the *amp* gene only [the polylinker and *Fl* origin was removed]. The blunt-ended and ligated *Acl*III and *Bsp*HI junction forms a unique *Ksp*I site. The 1732 bp plasmid was cut with *Ksp*I, blunt-end modified with *T4* polymerase, and a 20 bp *Rind*III-*K*incII fragment (blunt-end modified with the *Klenow* enzyme) obtained from the pUC19 polylinker was ligated into the blunt *Ksp*I site of the plasmid.

-34-

The *HindIII* site from the blunted polylinker was regenerated, and then positioned adjacent to the bacterial origin of replication. The resulting plasmid was then cut at the unique *PstI*/*Sse8387I* site, and an *Sse8387I*-*PvuII*-*Sse8387I* oligonucleotide, having the sequence: 5'-GGACGCTGCTTCA-3' (SEQ ID NO.: \_\_\_\_), was ligated therein. The remaining unique *BspHI* site was cut, blunt-end modified with *X*enow enzyme, and an *AscI* linker oligonucleotide, having the sequence: 5'-GAAGCGCGCTTC-3' (SEQ ID NO.: \_\_\_\_), was ligated therein, eliminating the *BspHI* site. The resulting plasmid was called pWec.

In order to create the pWadhlacZ construct, a CMVlacZ expression cassette (comprising a nucleotide sequence flanked 5' and 3' by *AAV* ITRs, containing the following elements: a CMV promoter, the *hcm1* 1st intron, an *adhIacZ* fragment and an SV40 early polyadenylation site) was inserted into the unique *PvuII* site of pWec using multiple steps such that the CMV promoter was arranged proximal to the bacterial *amp* gene of pWec.

More particularly, a CMVlacZ expression cassette was derived from the plasmid psub201CMV, which was constructed as follows. An oligonucleotide encoding the restriction enzyme sites:

*NotI*-*NruI*-*SnaBI*-*AgaI*-*BstBI*-*BstIII*-*KcoI*-*HpaI*-*BspEI*-*PstI*-*RsaI*-*NotI* and having the following nucleotide sequence:

5'-GCGGCGCAGCGTACGTACCGCTTCGAGCGCGACGCGGACCACTCTT  
AACTCCGACACTGCGACCGCGCGCG-3' (SEQ ID NO.: \_\_\_\_), was synthesized and cloned into the blunt-end modified *KasI*-*EatI* site (partial) of pUC119 to provide a 2757 bp vector fragment. A 653 bp *SpeI*-*SacII* fragment containing a nucleotide sequence encoding a CMV immediate early promoter was cloned into the *SnaBI* site of the 2757 bp vector fragment. Further, a 269

bp PCR-produced *BstBI*-*BstBI* fragment containing a nucleotide sequence encoding the *hcm1* 1st intron which was derived using the following primers:

5'-AAATTCGACACTGCGGACGACGAG-3' (SEQ ID NO.: \_\_\_\_)  
and 3'-aaattcgaaagtgtagccctttg-5' (SEQ ID NO.: \_\_\_\_), was cloned into the *BstBI* site of the 2757 bp vector fragment, and a 335 bp *HpaI*-*BamHI* (blunt-end modified) fragment containing the SV40 early polyadenylation site from the pCMV- $\beta$  plasmid (Clontech) was cloned into the *HpaI* site of the subject vector fragment. The resulting construct was then cut with *NotI* to provide a first CMV expression cassette.

Plasmid pW1909adhIacZ was constructed as follows. A 4723 bp *SpeI*-*EcoRV* fragment containing the *AAV* rep and *cap* encoding region was obtained from the plasmid pCM1909 (ATCC Accession Number 59871). The pCM1909 plasmid is a high efficiency *AAV* helper plasmid having *AAV* rep and *cap* genes with an *AAV* p5 promoter region that is arranged in the construct to be downstream from its normal position (in the wild type *AAV* genome) relative to the rep coding region. The 4723 bp fragment was blunt-end modified, and *AscI* linkers were ligated to the blunted ends. The resultant fragment was then ligated into the unique *AscI* site of pWadhlacZ and oriented such that the *AAV* coding sequences were arranged proximal to the bacterial origin of replication in the construct.

Plasmid pW1909adhIacZ includes the bacterial beta-galactosidase ( $\beta$ -gal) gene under the transcriptional control of the cytomegalovirus immediate early promoter (CMVIE).

#### B. Construction of pW1909EPO.

Plasmid pW1909adhIacZ was modified to express human erythropoietin (EPO) by replacing the *adhIacZ* gene with a 718 base pair *PvuII*-*NcoI* fragment

of human EPO cDNA (Wen et al. (1993) *Blood* 5:1507-1516) and by cloning a 2181 bp *Cla*I-*Eco*RI *lacZ* spacer fragment (noncoding) into the *Pml*I site of the vector.

### 5 C. Construction of pAAV-GAA.

A plasmid containing the human lysosomal enzyme acid  $\alpha$ -glucosidase (GAA) coding region was constructed as follows. A 3.2 kb cDNA clone containing the coding sequence for human GAA beginning 207 bps downstream from the initiation codon (GenBank Accession Numbers: M34424 and Y00839; Martinik et al. (1990) *DN*A Cell Biol. 2:85-94) was cloned into the *Eco*RI site of Bluescript KS (Stratagene). Additional 5' sequence was generated using polymerase chain reaction (PCR) with reverse-transcribed poly-A mRNA (obtained from normal human fibroblasts) as the template. The 5' primer was constructed with a *Xba*I restriction site and bps -1 to 12 of the published sequence (Martinik et al. (1986) *Proc. Natl. Acad. Sci. USA* 83:9641-9644; Hofeise et al. (1983) *Eur. Mol. Biol. Organ.* 2:1657-1704). The 3' primer was synthesized from basepairs 1001 to 1018. Using *Xba*I and the unique internal *Stu*I site at position 776, the PCR product was ligated to the partial cDNA to form the full-length GAA-encoding plasmid. The full length cDNA was truncated at a unique *Sph*I restriction site, and cloned into the expression vector p1.1c to result in the pAAV-GAA construct having the GAA coding region under transcriptional control of the CMV-IE promoter.

The p1.1c expression vector was constructed as follows. pUC119 was partially digested with *R*saI and *E*arI, and a 2713 bp vector fragment containing the ampicillin resistance gene, the *col*1 origin of replication and the M13 origin of replication, was isolated, blunt end modified, and ligated to a synthetic DNA polylinker encoding the restriction enzyme sites *Not*I-*H*uI-*S*naBI-*X*baI-*S*fuI-*B*ssHII-*S*agI-

*N*coI-*P*meI-*B*spRI-*P*mlI-*R*ecII-*N*otI, and having the following nucleotide sequence:

5'-CCGCCGCCACCGCTTGTAAACACCGGTTGAAACCGG  
CAGCGCCGACATGCTTAACTCCGACACGACGACGACGACGACG-  
3' (SEQ ID NO.: \_\_\_\_). The ligation was conducted such that the *H*uI end of the polylinker was ligated to the *R*saI site of the plasmid. A 653 bp *S*paI-*S*acII fragment encoding the CMV immediate-early promoter, a 269 bp PCR-produced *S*fuI-*S*fuI produced fragment encoding the hGH 1st intron (derived using the following primers: 5'-AAATTCGAACAGTAACGCCCTTG-3' (SEQ ID NO.: \_\_\_\_)) and 3'-AAATTCGAACCTGGGACAAACCGAG-5' (SEQ ID NO.: \_\_\_\_)), a 183 bp *B*ssHII-*B*ssHII polylinker fragment from Bluescript II SK-, and a 135 bp *H*paI-*B*ssHII (blunted) fragment containing the SV40 early polyadenylation site from pCMV- $\beta$  (Stratagene), were cloned into the *S*naBI, *S*fuI, *B*ssHII, and *P*meI sites, respectively, of the aforementioned plasmid. The orientation of the polylinker relative to the intron and polyadenylation site was intron-polylinker (5'-*S*acI-3'-*X*baI)-polyadenylation site. The polylinker was further modified by removing the 86 bp *S*acI-*X*baI polylinker fragment and replacing it with the following synthetic *S*acI to *X*baI fragment encoding the restriction enzyme sites *S*acI-*C*laI-*E*coRI-*S*maI-*B*ssHII-*X*baI-*S*aiI-*P*stI-*B*stXI-*E*coRV-*B*stXI-*o*loneganuclease-*H*indIII-*X*baI, having the following nucleotide sequence:

5'-GAGCTCATCGATTGAATTCGCCGGGATCTCTTAAGTGCACCTGCAACC  
ACTGTGTGATATACAAACACACTGCTAGGATTAACAGGTAATTCGAG-3'  
(SEQ ID NO.: \_\_\_\_).

### Virus and Cell Lines

Adenovirus type 2 (Ad2), available from the American Type Culture Collection, ATCC, catalogue number VR846, was used as helper virus to encapsidate vectors.

The human 293 cell line (Graham et al. (1977) *J. Gen. Virol.* 26:59-77, available from the ATCC under Accession no. CRL1573), which has adenovirus E1a and E1b genes stably integrated in its genome, was cultured in complete Dulbecco's modified Eagle's media (DMEM; Bio-Whittaker, Walkersville, MD) containing 4.5 g/L glucose, 10% heat-inactivated fetal bovine serum (FBS; Hyclone, Logan, Utah), 2 mM glutamine, and 50 units/mL penicillin and 50 µg/mL streptomycin.

The C2C12 murine myoblast cell line, available from the ATCC, Catalogue Number CRL1772, was cultured in DMEM with 20% fetal calf serum (FCS), 1% chick embryo extract and 5 µg/mL gentamicin.

Fetal human skeletal myoblasts (clonatics) were cultured in Ham's F-12 human growth medium, containing 20% FCS and 5 µg/mL gentamicin.

The above cell lines were incubated at 37 °C in 5% CO<sub>2</sub>, and were routinely tested and found free of mycoplasma contamination.

#### Production of Recombinant AAV Virions

Recombinant AAV virions were produced in human 293 cells as follows. Subconfluent 293 cells were cotransfected by standard calcium phosphate precipitation (Miyeler et al. (1979) *Proc. Natl. Acad. Sci USA* 76:11373-11376) with one of the AAV vector/helper plasmid constructs, pM1909adhlacZ or pM1909EPO, or with pAAV-GA and the pM1903 helper plasmid. After 6 hours, the transfected cells were infected with Ad2 in fresh medium at a multiplicity of infection (MOI) of 2, and incubated at 37 °C in 5% CO<sub>2</sub> for 70 hours prior to harvest. Pelleted cells were lysed in Tris buffer (10 mM Tris, 150 mM NaCl, pH 8.0) by three cycles of freeze-thaw. The lysate was clarified of cell debris by centrifugation at 12,000 X g, and the crude-cell lysate was layered onto a cesium

-39-

chloride cushion for isopycnic gradient centrifugation. Recombinant AAV virions (rAAV-lacZ, rAAV-EPO, or rAAV-GA virions) were extracted from the resulting gradient by isolating the fractions with an average density of approximately 1.38 g/mL, resuspended in Hepes buffered saline (HBS) containing 50 mM Hepes (pH 7.4) and 150 mM NaCl. The preparations were then heated at 56 °C for approximately 1 hour to inactivate Ad2.

#### Assay of rAAV by Dot-blot Hybridization

Recombinant AAV virions were DNase I digested, proteinase K treated, phenol-chloroform extracted, and DNA precipitated with sodium acetate-glycogen (final concentrations = 0.3 M sodium acetate and 160 µg/mL, respectively). DNA samples were denatured (200 µL of 2X alkaline solution (0.8 M NaOH, 20 mM DTTA) added to DNA sample) for 10 minutes, then added to appropriate wells in a dot-blot apparatus, and blotted onto wet zeta probe membrane (Elopad), by applying suction until wells were empty. Then, 100 µL of 1X alkaline solution was added; after 5 minutes, wells were emptied by suction. The membrane was rinsed in 2X SSC (Sambrook et al., *supra*) for 1 min, drained, air dried on filter paper, then baked in vacuum at 80 °C for 30 min. The membrane was then prehybridized for 30 min at 65 °C with 10 mL hybridization buffer (1% SDS, 0.25 M sodium phosphate, pH 7.2, 1 mM EDTA). Buffer was replaced with 10 mL fresh solution, freshly boiled probe added, and hybridized overnight at 65 °C. The membrane was washed twice with 25 mL of wash-1 buffer (5% SDS, 40 mM sodium phosphate, pH 7.2, 1 mM EDTA) for 20 min at 65 °C and twice with wash-2 buffer (1% SDS, 40 mM sodium phosphate, pH 7.2, 1 mM EDTA). The membrane was wrapped in plastic film, exposed to radiographic film, and appropriate dots excised from the membrane

-40-

to determine radioactivity by scintillation counting, and quantitated by comparison with standards. Titers of rAAV virion were routinely in the range of approximately  $10^{12}$  genomes/mL.

#### Assay for Contaminating Helper Adenovirus

Contaminating infectious adenovirus was assayed as follows. Samples from the purified rAAV virion stocks were added to 50% confluent 293 cells (cultured in 12 well dishes at  $1 \times 10^5$  cells/well), and the cultures were passaged for 30 days (e.g., the cultures were split 1 to 5, every 3 days) or until the culture exhibited 100% cytopathic effect (CPE) due to adenovirus infection. Cultures were examined daily for CPE, and the day upon which each experimental culture showed 100% CPE was noted. Reference 293 cell cultures infected with a range of known amounts of adenovirus type-2 (from 0 to  $1 \times 10^3$  plaque forming units (pfu)/culture) were also prepared and treated in the same manner. A standard curve was then prepared from the data obtained from the reference cultures, where the adenovirus pfu number was plotted against the day of 100% CPE. The titer of infectious adenovirus type-2 in each experimental culture was then readily obtained as determined from the standard curve. The limit of detection of the assay was 100 pfu/mL. The presence of wild-type AAV contamination, analyzed by dot-blot hybridization, was approximately 7 logs lower than recombinant virion concentration.

#### Differentiation of Myoblasts

C2C12 myoblasts were transduced either while actively dividing, or as a differentiated cell culture. Differentiation was induced by placing subconfluent myoblasts in curine differentiation medium (DMEM containing 2% horse serum and standard

concentrations of glutamine and penicillin-streptomycin) for an interval of four days prior to transduction in order to induce myoblast fusion and formation of differentiated myotubes.

Recal human skeletal myoblasts were differentiated in human differentiation medium (DMEM containing 10% horse serum and 5 µg/mL gentamicin). Verification of differentiation was performed by microscopic analysis to determine the presence of multinucleated myotubes in culture.

#### Example 1

##### Expression of rAAV-LacZ in Terminally Differentiated Adult Rat Cardiomyocytes

The ability of recombinant AAV virions to transduce terminally differentiated adult cardiomyocytes was established *in vitro*. Cardiomyocytes were harvested by coronary perfusion with collagenase of adult rat hearts (Fischer 344, Harlan Sprague Dawley, Indianapolis, IN). Cardiomyocytes were grown on laminin-coated glass coverslips and exposed to rAAV-LacZ virions for 4 hours. After 72 hours, the cells were stained for  $\beta$ -galactosidase activity. AAV expression was detected by blue staining of the bisected cells. These studies demonstrated the ability of AAV virions to transduce terminally differentiated cells. The transduction efficiency *in vitro* was 30% of adult cells at a multiplicity of infection (MOI) of  $10^4$  genomes per cell.

#### Example 2

##### Stability of rAAV-LacZ Expression *In Vivo*

Adult Fischer rats were used to analyze expression of transgenes *in vivo*. Incremental doses of rAAV-LacZ virions were injected into the left ventricular apex of the heart using either a

subxyphoid or lateral thoracotomy approach. Mice particularly, experimental animals were anesthetized with Metofane followed by a subxyphoid incision to expose the diaphragmatic surface of the heart. Apical cardiac injections were performed with a glass micropipette. Recombinant virion was diluted in normal saline and injected at a volume of 20-50µL.

At varying times post-injection, hearts were harvested and examined for  $\beta$ -galactosidase production and for the presence of infiltrating mononuclear cells. For 5-Bromo-4-chloro-3-indolyl  $\beta$ -D-galactoside histochemical determination, frozen sections (6µm) were fixed in 0.5% glutaraldehyde and stained for  $\beta$ -galactosidase activity as described (Sanas et al. (1986) "Use of Recombinant Retrovirus to Study Post-Implantation Cell Lineage in Mouse Embryos," *EMBO J* 5:3133-3142). Paraffin sections (5µm) were stained with hematoxylin/eosin. Sections were examined for infiltrating mononuclear cells.

The above-described histochemical studies showed greater than 50% transduction of cardiomyocytes in the region of injection at each time point examined. Further, there was no inflammatory cell infiltrate noted during the course of analysis.  $\beta$ -galactosidase staining was observed to persist in cardiac muscle for at least two months following gene transfer.

#### Example 2

##### In vivo Transduction of Murine Skeletal Muscle using rAAV-LacZ Virions

Recombinant AAV-LacZ virions were injected into muscle tissue of mice, and transduction assessed by  $\beta$ -gal activity. Particularly, *in vivo* transduction was performed by intramuscular (IM) injection of recombinant AAV virions into the skeletal muscle of healthy 6-8 week old Balb/c mice (Jackson

laboratories, Bar Harbor, ME, Simonsen Laboratories, Gilroy, CA, or Harlan Laboratories) under either Metofane (Pitman-Moore, Mundelein, IL) or ketamine-xylazine anesthesia. The mid-portion of each tibialis anterior muscle was exposed via a 1 cm incision.

Injections into the tibialis anterior were carried out using a micro-capillary tube attached to a Hamilton syringe to administer the following formulations at a depth of 2 mm: phosphate-buffered saline (PBS) alone (negative control); or PBS containing either rAAV-LacZ virions or pM1909adhlacZ plasmids.

Tissue samples of tibialis anterior muscle, forelimb muscle, heart, brain and liver were obtained for analysis of  $\beta$ -galactosidase expression. One tibialis anterior muscle from each animal was processed for cross-sectional  $\beta$ -galactosidase analysis, and total  $\beta$ -galactosidase was determined from a crude homogenate of the other muscle using a chemiluminescent assay.

For histochemical detection of  $\beta$ -galactosidase, muscle samples were snap-frozen in dry ice-cooled isopentane, followed by serial transverse sectioning (10 µm) and processing according to previously described methods (Sanas et al. (1986) *EMBO J* 5:3133-3142). The cross-sectional area of the tibialis anterior expressing  $\beta$ -galactosidase was determined as follows: after counter-staining with nuclear fast red (Vector Labs), the X-gal stained tissue was digitally photographed and the cross-sectional area of stored images was determined using NIH Image software.

The GALACTO-LIGHT™ (Tropix, Bedford, MA) chemiluminescent reporter assay kit was used to detect total  $\beta$ -galactosidase activity in the entire tibialis anterior muscle. Standard curves were prepared from known amounts of purified  $\beta$ -galactosidase (Sigma, St. Louis, MO) resuspended in non-transduced muscle

homogenate.  $\beta$ -galactosidase activity is expressed as either: nanograms of  $\beta$ -galactosidase, normalized for the entire muscle, minus background activity; or in terms of relative light units (RLU) as quantified by luminometer. Forelimb muscle, cardiac muscle, brain tissue and liver samples were assayed in an identical fashion.

#### A. Time course of $\beta$ -galactosidase

##### Expression.

A single intramuscular injection into the left and right tibialis anterior muscles (under direct vision) was used to deliver  $8 \times 10^9$  rAAV-LacZ in a PBS vehicle. Four animals were injected with PBS alone. Animals were sacrificed at 2, 4, 8, 12, 24 and 32 weeks after injection, and the tibialis anterior muscle was excised and analyzed for the presence of bacterial  $\beta$ -galactosidase ( $n=5$  for each group) as described above.

Efficiency of the LacZ gene transfer was assessed by cross-sectional tissue staining and chondrofluorescent assay. As can be seen in Figure 1 and in Table I below, gene expression persisted for at least 32 weeks. In addition, two weeks after injection of the recombinant virions, 14% of the muscle cross-sectional area expressed  $\beta$ -galactosidase, while at 32 weeks, 24% of the muscle cross-sectional area expressed  $\beta$ -galactosidase. Negative control tibialis anterior muscle (obtained from the animals injected with PBS alone) showed no background staining. Muscle  $\beta$ -galactosidase activity was also determined in the contralateral injected muscle. This study also revealed persistent expression for at least 32 weeks, in agreement with the cross-sectional fiber analysis (Table I). Further, the observed  $\beta$ -galactosidase was sustained with minimal inflammatory cell infiltrate. These data demonstrate that rAAV-

LacZ virion administration into muscle results in stable expression of the transgene for at least 8 months.

Upon histological examination, positive staining filled the cytoplasm of the transduced myofibers and was observed through large contiguous portions of the muscle. Serial transverse sections revealed that blue staining extended throughout the length of the muscle fiber. Diffraction-interference contrast microscopy revealed a clear delineation between positively and negatively stained myofibers (Figure 2), suggesting that recombinant virion delivery was limited by structural barriers such as epimyseal or perimyseal connective tissue. Homogenates prepared from brain, heart, liver, and forelimb muscle displayed no  $\beta$ -galactosidase activity when compared with the background activity of the negative control animals.



Table 1  
Time Course of  $\beta$ -galactosidase Expression  
Following Single Injection with rAAV-LacZ.

Time (weeks)	$\beta$ -galactosidase expression (ng/muscle)	Percent cross- sectional area expressing $\beta$ - galactosidase (%)
2	1441 $\pm$ 453	18 $\pm$ 6
4	951 $\pm$ 176	20 $\pm$ 4
8	839 $\pm$ 436	24 $\pm$ 3
12	1874 $\pm$ 521	29 $\pm$ 5
24	2579 $\pm$ 1165	22 $\pm$ 3
32	1242 $\pm$ 484	24 $\pm$ 3

The left and right tibialis anterior muscles were injected with  $8 \times 10^8$  rAAV-LacZ. One number of the pair of injected muscles was processed for  $\beta$ -galactosidase expression ( $n=3 \pm$  SEM). The other muscle was processed for histochemical detection of  $\beta$ -galactosidase and determination of the cross-sectional area of the tibialis anterior expressing  $\beta$ -galactosidase. Mean cross-sectional areas  $\pm$  SEM are shown.

#### B. Dose-response assay.

To determine the effective dose range for rAAV-LacZ in vivo, recombinant virions were injected into the tibialis anterior muscle of 6-8 week old Balb/c mice, and transduction assessed by  $\beta$ -galactosidase activity as measured by ELACTO-LIGHT<sup>TM</sup> relative light units (RLU). As can be seen in Figure 3, at two weeks post-injection, the observed RLU ranged from approximately  $0.2 \times 10^3$  RLU/muscle (injected with  $8 \times 10^8$  rAAV-LacZ) to approximately  $1.1 \times 10^3$  RLU/muscle (injected with  $1.6 \times 10^{11}$  rAAV-LacZ).

The levels of expression of  $\beta$ -galactosidase measured in RLU correspond to the percentage of  $\beta$ -galactosidase-positive muscle fibers on cross-sectional analysis. For example,  $0.2 \times 10^3$  RLU corresponds to approximately 1%  $\beta$ -galactosidase

positive muscle fibers and  $1.1 \times 10^3$  RLU corresponds to approximately 60%  $\beta$ -galactosidase positive muscle fibers.

#### C. Comparison of $\beta$ -galactosidase expression efficiency.

A comparison of  $\beta$ -galactosidase expression efficiency obtained by in vivo transduction of mice using either rAAV-LacZ virions, or plasmid DNA containing the same lacZ expression cassette (pR1909adhLacZ) was carried out as follows. Either  $8 \times 10^8$  rAAV-LacZ, or 100  $\mu$ g of pR1909adhLacZ, was injected into the tibialis anterior muscle of 6-8 week old Balb/c mice. Two weeks post-injection,  $\beta$ -galactosidase activity was assessed using the ELACTO-LIGHT<sup>TM</sup> chemiluminescent reporter assay kit, as described above. Administration of the recombinant virions resulted in 1441 ng  $\beta$ -galactosidase/muscle ( $n=5$ ), while administration of 100  $\mu$ g of the plasmid DNA, a typical in vivo plasmid DNA dosage (Whalen et al. (1995) Hum. Gene Ther. 5:151-159), resulted in 12 ng  $\beta$ -galactosidase/muscle ( $n=4$ ). This dosage of pR1909adhLacZ plasmid DNA is equivalent to  $2.2 \times 10^{13}$  single stranded genomes, demonstrating that gene delivery by the recombinant virions was substantially more efficient than delivery of an equal molar quantity of vector DNA.

Example 4In vitro Transduction of Murine  
Myotubes and Myoblasts

5 cultured muscle cells are appropriate targets for  
recombinant AAV virion transduction, and to assess the  
ability of such cells to express a transduced gene,  
the following study was carried out. Murine C2C12  
cells were selected since these cells have been  
10 extensively studied as a model for mammalian  
myogenesis (Blau et al. (1993) Trends Genet. 2:269-  
274), and can be induced to differentiate by growth in  
reduced serum medium.

15 In the study, C2C12 myoblasts (dividing  
cells) were seeded in cell culture plates at a density  
of  $2 \times 10^4$  cells/cm<sup>2</sup>, maintained in growth media (GM)  
until confluent, split, and then either cultured in GM  
or cultured for 5 days in murine DM. Differentiation  
was verified by the microscopic presence of  
20 multinucleate myotubes, representing fused myoblasts  
(differentiated C2C12 cells).

The C2C12 myotubes and myoblasts were  
transduced in culture with purified rAAV-hEPO virions  
at a MOI of  $10^5$  in OptiMEM (Gibco BRL). In the  
25 myotube cultures, DM was added after virion  
adsorption. The culture media of the transduced cells  
was changed 24 hours prior to collection of  
supernatants at 3, 8 and 14 days following  
transduction. Secretion of hEPO was assessed by ELISA  
30 using the human erythropoietin Quantikine R&D kit  
(available from R and D Systems, Minneapolis, MN)  
according to manufacturer's recommendations.

35 The results of the study show that hEPO is  
secreted from both the transduced myotubes and  
myoblasts. The levels of hEPO secretion increased in  
the myotubes over the first seven days post-  
transduction (Figure 4). As can be seen by reference

-49-

to Figure 5, a dose-dependent increase in the  
secretion of hEPO was also observed in the transduced  
C2C12 myotubes. Eight days post-transduction of the  
5 myotubes, hEPO levels peaked at  $>3400$  uU/mL. These  
data demonstrate that transduction with rAAV-hEPO of  
both myotubes or myoblasts results in hEPO secretion  
by the transduced cells, and that in short-term  
myotube cultures, hEPO is synthesized and secreted in  
a dose-dependent manner.

Example 5In vitro Transduction of Human  
Myotubes Using rAAV-hEPO Virions

15 To determine if differentiated primary human  
muscle cells are able to express hEPO following  
transduction with rAAV-hEPO, the following study was  
carried out. Primary fetal human skeletal myoblasts  
were seeded in cell culture plates at a density of  $2 \times$   
20  $10^4$  cells/cm<sup>2</sup>, grown to confluence in appropriate  
growth media, and then cultured for 14 days in human  
DM. Differentiation was verified by microscopic  
examination for multinucleate cells. In vitro  
transduction was carried out by adding purified rAAV-  
25 hEPO virions to the cultured myotubes in OptiMEM  
medium (Gibco BRL). DM was added to the cultures  
after virion adsorption. Control cultures were  
transduced with rAAV-LacZ.

Culture media was changed 24 hours prior to  
collection of supernatants at day 3, 8 and 14 post  
30 transduction. Secreted hEPO levels were assayed by  
ELISA as described above in Example 4.

35 As can be seen in Figure 6, the transduced  
human myotubes secreted hEPO into the culture in a  
dose-dependent manner. No detectable hEPO activity was  
measured in the control cultures. Secretion of hEPO  
increased over the 14-day interval post-transduction.  
These data demonstrate that primary human myotubes

-50-

transduced by recombinant rAAV virions are capable of expressing and secreting erythropoietin.

#### Example 6

##### Systemic Delivery of Human Erythropoietin *In Vivo* by Intramuscular Administration of rAAV-HEPO

Recombinant rAAV virions encoding hEPO were administered to adult healthy Balb/c mice *in vivo* to determine if a systemic level of hEPO can be produced, and a biological response obtained. At various time points after administration, blood was obtained from the orbital venous plexus under anesthesia. Serum hEPO levels were determined by ELISA as described above. Red cell counts were done by hemocytometer, hematocrit was determined by centrifugation of blood in micro-capillary tubes, and hemoglobin concentration was analyzed by cyanmethemoglobin assay (DHA, Arlington, TX) according to manufacturer's specifications and compared with a standard (Scanbio Laboratory, San Antonio, TX) analyzed at 570 nm on a spectrophotometer. Reticulocytes were analyzed by either new methylene blue stain, or by FACS analysis of chlorole orange stained peripheral blood samples (RETIC-COUNT, Becton-Dickinson, Mountain View, CA); the results of data obtained by either of these methods were similar.

An initial experiment revealed that high levels of hEPO and elevated hematocrits were maintained for >100 days in mice injected IM with  $6.5 \times 10^{11}$  rAAV-HEPO. Next, adult female Balb/c mice were injected IM in both hind limbs with a single administration of rAAV-HEPO at dosages ranging from  $3 \times 10^9$  to  $3 \times 10^{11}$  particles. Control animals were injected with rAAV-lacZ. The resulting serum hEPO levels were analyzed and are reported below in Table IJ. As can be seen, a well-defined dose-response was obtained 20, 41, 62 and 83 days post injection.

-51-

The time course of hEPO secretion by animals receiving rAAV-HEPO is depicted in Figure 7. As can be seen, serum levels of hEPO increased with time to plateau at from 6 to 8 weeks after injection.

The biological activity of secreted hEPO can be monitored by elevation of hematocrit in the experimental animals. A comparison of circulating hEPO levels versus hematocrit is shown in Table II. The comparison shows that hematocrit increased with time and increasing recombinant virion dose. Further, stable elevation in hematocrit has been observed for up to 40 weeks in a group of experimental animals injected with rAAV-HEPO. Control animals had undetectable levels of hEPO ( $<2.5$  mU/mL, the lower limit of detection for the assay).

These results indicate that persistent and stable high-level secretion of hEPO, with a corresponding elevation in hematocrit, is established following a single IM administration of rAAV-HEPO.

In addition, comparison of the expression of hEPO by animals injected IM with rAAV-HEPO ( $3 \times 10^{11}$  single-stranded genomes) and animals injected IM with the pM1905EPO plasmid ( $1.4 \times 10^{13}$  double-stranded genomes in 100 $\mu$ g DNA) shows that the recombinant virions gave rise to significantly greater levels of hEPO expression. As reported in Table II, 20 days post-injection, recombinant virion-injected animals had serum levels of  $445 \pm 98$  mU/mL, while the plasmid-injected animals had levels of  $8 \pm 10$  mU/mL. At 41 days post-injection, levels in the recombinant virion-treated animals had risen to  $725 \pm 112$  mU/mL, while the levels in the plasmid-treated animals had dropped below the level of detection. The animals receiving rAAV-HEPO exhibited approximately 60-fold more circulating hEPO with 100-fold less input genomes at 20 days post-injection, or approximately 5000-fold greater secretion per genome. At 41 days post-

-52-

Injection, this difference was even greater, since the plasmid expression was below the level of detection.

Table II EPO Expression and Hematocrit: rAAV-hEPO Dose-Response								
Dose	Days after Administration							
	20 days		41 days		62 days		83 days	
	EPO	HCT	EPO	HCT	EPO	HCT	EPO	HCT
$3 \times 10^{11}$	445±98	74.2±1.2	725±112	82.3±1.2	769±61	86.5±1.4	723±253	88.3±0.7
$1 \times 10^{11}$	85±14	72.8±1.5	212±23	79.5±1.7	234±75	83.2±0.2	220±51	83.2±2
$3 \times 10^{10}$	17±5	60.0±3.5	34±17	74.7±1.2	55±28	78.7±2.0	73±45	80.0±3
$1 \times 10^{10}$	3±1	52.9±1.8	11±3	61.5±1.9	12±8	68.4±4.6	15±5	70.8±8
$3 \times 10^9$	<2.5	49.9±1.4	<2.5	53.5±2.5	<2.5	57.0±2.4	4±4	57.5±3
i.v.	7±3	54.7±3.2	13±2.0	<64.4±5.3	10.1±0.7	70.8±8	21±10	74.6±7
Control	<2.5	48.9±1.0	<2.5	49.1±0.8	<2.5	48.1±0.7	<2.5	48.2±9
Plasmid	8±10	50±3.0	<2.5	50.2±1.0	<2.5	47.8±0.9	N.D.	N.D.

Values representing means ± standard deviation (SD). EPO = serum levels of human EPO (mU/mL) in Balb/c mice; HCT = hematocrit (%); N.D. = not done; i.v. = intravenous injection with  $3 \times 10^{11}$  rAAV-hEPO; Plasmid = injection with 100 µg plasmid DNA ( $1.4 \times 10^{13}$  double-stranded plasmid molecules); Control = injection with  $3 \times 10^{11}$  particles of rAAV-lacZ.

Example 7A. Comparison of hEPO Secretion from rAAV-hEPO Administered by IM or IV Routes

A comparison of the circulating levels of hEPO resulting from IM and IV routes of administration was analyzed to determine which method of gene delivery results in higher levels of systemic hEPO. Balb/c mice were injected with  $3 \times 10^{11}$  rAAV-hEPO using either the IM route as described above, or intravenously (IV) in PBS in a total volume of 50  $\mu$ L via the lateral tail vein. Serum hEPO levels were determined by ELISA using the methods described above.

As shown in Table II, hEPO levels resulting from the IV administrations were significantly lower than the group that received the virions by the IM route. In particular, at 20 days post-injection, the IM route resulted in levels of hEPO of  $445 \pm 38$  nU/mL, while the IV route produced  $7 \pm 3.0$  nU/mL. At 41 days post-injection, the hEPO level observed with the IM route was  $725 \pm 112$  as compared with  $13 \pm 2.0$  nU/mL by IV, or approximately 60-fold more efficacious. These data demonstrate that the IM route of injection resulted in higher systemic levels of hEPO, and suggest that interstitial delivery in muscle results in improved transduction by the recombinant AAV virions.

Example 8In Vitro and In Vivo Transduction of Muscle Cells Using rAAV-GAA Virions

Cardiomyopathy in infancy is frequently due to inherited metabolic disease. One such metabolic disease, glycogen storage disease type II (Pompe's disease) is an inherited cardiomyopathy caused by a deficiency in the lysosomal enzyme, acid  $\alpha$ -glucosidase (GAA). GAA functions to cleave  $\alpha$ -1,4 and  $\alpha$ -1,6 linkages of lysosomal glycogen to release

monosaccharides. Loss of enzyme activity results in accumulation of lysosomal glycogen in striated muscle, and is characterized by lysosomal rupture, contractile apparatus disruption and glycogen infiltration. Currently, no effective treatment is available.

Accordingly, the following studies were carried out to determine whether the recombinant AAV virions of the present invention can be used to obtain long-term expression of GAA in transduced muscle cells.

A. In Vitro Transduction of Human Skeletal Muscle Cells with rAAV-hGAA Virions.

Human skeletal myoblasts were seeded in cell culture plates at a density of  $2 \times 10^4$  cells/cm<sup>2</sup>, grown to confluence in appropriate growth media, and then cultured for 14 days in human DM. Differentiation was verified by microscopic examination for multinucleate cells. In vitro transduction was carried out by adding purified rAAV-hGAA virions at an MOI of  $2 \times 10^3$  to the cultured myotubes in OptiMEM medium (Gibco BRL). DM was added to the cultures after virion adsorption. Transduced control cultures were established by transducing the myotubes with rAAV-lacZ as the same MOI, and negative controls were established by culturing non-transduced myotubes.

Culture media was changed 24 hours prior to collection of supernatants at day 3, 8 and 14 post transduction. hGAA expression was determined by enzymatic assay. Specifically, cell monolayers were harvested with 0.05% trypsin in Puck's saline A containing 0.02% ethylenediaminetetraacetic acid. After quenching the trypsin with growth media, cells were centrifuged, the cell pellet washed with PBS, and resuspended in distilled water. Following three freeze-thaw cycles, the samples were microcloned at  $10,000 \times g$ . Protein in the supernatant was determined

using the Bicinchoninic acid method (Pierce) with bovine serum albumin (BSA) as the standard. GAA activity was measured using cleavage of the glycogen analog 4-methylumbelliferyl- $\alpha$ -D-glucoside, by a modification of a previously described method (Galjaard et al. (1973) *Clin. Chim. Acta.* 49:361-375; Galjaard, H. (1973) *Pediatr. Res.* 2:56). Assays contained 30  $\mu$ g protein in 125  $\mu$ L water. Two volumes of 200 mM sodium acetate (pH 4.3) and 750 mM 4-methylumbelliferyl- $\alpha$ -D-glucoside (from a stock solution of 200 mM in dimethylsulfoxide) were added and the samples incubated at 37°C for one hour. Reactions were stopped with 625  $\mu$ L sodium carbonate (pH 10.7). Once cleared and alkalized, the 4-methylumbelliferyl fluorosces. Measurements were made with a fluorescence spectrophotometer with excitation at 365 nm and emission at 448 nm. Assay measurements were compared against 4-methylumbelliferone (Sigma, St. Louis, MO) standards. Zero protein and zero time blanks were used.

The observed GAA activity is reported in Figure 8 which demonstrates that in vitro transduction of the human myotubes results in high level GAA expression at day 8 and 14 post transduction.

#### B. In vivo transduction of skeletal muscle using rAAV-GAA virions.

Recombinant AAV virions encoding human GAA were administered to adult Balb/c mice in vivo to determine if systemic levels of GAA can be produced by the transduced cells. Muscle tissue was isolated at various time points after administration, and processed for GAA activity using an enzymatic assay.

In the study, tibialis anterior muscle in Balb/c mice was surgically exposed, and a single intramuscular injection into the left and right muscles (under direct vision) was used to administer

the following formulations: phosphate-buffered saline (PBS) alone (negative control); or PBS containing either  $2 \times 10^{10}$  rAAV-hGAA or rAAV-LacZ (for a total of  $4 \times 10^{10}$  virions/animal).

Tissue samples of tibialis anterior muscle were obtained at 1, 4 and 10 weeks following transduction. The tissue samples were prepared by homogenization in water followed by three freeze-thaw cycles. Freeze-thaw lysates were microcentrifuged, and the resultant supernatant assayed for GAA activity as described above. The results of the study are depicted in Figure 9. As can be seen, stable expression of GAA in the transduced mouse muscle cells was observed for ten weeks, demonstrating that the recombinant AAV virions of the present invention are able to establish efficient expression of a functional lysosomal protein in transduced cells, and thus provide a therapeutic approach for the treatment of glycogen storage disease.

Accordingly, novel methods for transferring genes to muscle cells have been described. Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without departing from the spirit and the scope of the invention as defined by the appended claims.

Deposits of Strains Useful In Practicing the Invention

A deposit of biologically pure cultures of the following strain was made with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland, under the provisions of the Budapest Treaty. The accession number indicated was assigned after successful viability testing, and the requisite fees were paid. Access to said cultures will be available during pendency of the patent application to one determined by the Commissioner to be entitled thereto under 37 CFR 1.14 and 35 USC 122. All restriction on availability of said cultures to the public will be irrevocably removed upon the granting of a patent based upon the application. Moreover, the designated deposits will be maintained for a period of thirty (30) years from the date of deposit, or for five (5) years after the last request for the deposit; or for the enforceable life of the U.S. patent, whichever is longer. Should a culture become nonviable or be inadvertently destroyed, or, in the case of plasmid-containing strains, lose its plasmid, it will be replaced with a viable culture(s) of the same taxonomic description.

This deposit is provided merely as a convenience to those of skill in the art, and is not an admission that a deposit is required. A license may be required to make, use, or sell the deposited materials, and no such license is hereby granted.

Strain Deposit Date ATCC No.  
PGN1909 July 20, 1995 69871

Claims:

1. A method for producing a composition useful for delivering a selected gene to a muscle cell or tissue, comprising:
  - (a) providing a recombinant adeno-associated virus (AAV) virion which comprises an AAV vector having said selected gene operably linked to control elements capable of directing the *in vivo* transcription and translation thereof; and
  - (b) combining the recombinant AAV virion with a pharmaceutically acceptable vehicle.
2. The method of claim 1, characterized in that the composition is useful for delivering a selected gene to a muscle cell or tissue derived from skeletal muscle.
3. The method of claim 2, characterized in that the muscle cell is a skeletal myoblast or a skeletal myocyte.
4. The method of claim 1, characterized in that the composition is useful for delivering a selected gene to a muscle cell or tissue derived from smooth muscle.
5. The method of claim 1, characterized in that the composition is useful for delivering a selected gene to a muscle cell or tissue derived from cardiac muscle.
6. The method of claim 5, characterized in that the muscle cell is a cardiomyocyte.

7. The method of any one of claims 1-6, characterized in that the selected gene encodes a therapeutic protein.

8. The method of claim 7, characterized in that the therapeutic protein is acid  $\alpha$ -glucosidase.

9. The method of any one of claims 1-8, characterized in that the control elements comprise a muscle-specific promoter sequence.

10. The method of any one of claims 1-8, characterized in that the control elements comprise an inducible promoter sequence.

11. A muscle cell or tissue transduced in vitro with a recombinant AAV virion, said virion comprising an AAV vector having a selected gene operably linked to control elements capable of directing the in vivo transcription and translation of said selected gene.

12. The transduced muscle cell of claim 11, characterized in that the muscle cell is a skeletal myoblast or a skeletal myocyte.

13. The transduced muscle cell of claim 11, characterized in that the muscle cell is a cardiomyocyte.

14. The transduced muscle cell of any one of claims 11-13, characterized in that the selected gene encodes a therapeutic protein.

15. The transduced muscle cell of claim 14, characterized in that the therapeutic protein is acid  $\alpha$ -glucosidase.

16. The transduced muscle cell of any one of claims 11-15, characterized in that the control elements comprise a muscle-specific promoter sequence.

17. The transduced muscle cell of any one of claims 11-15, characterized in that the control elements comprise an inducible promoter sequence.

18. A method of transducing a mammalian muscle cell in vitro, comprising:

(a) providing a recombinant AAV virion which comprises an AAV vector having a selected gene operably linked to control elements capable of directing the in vivo transcription and translation of said selected gene; and

(b) introducing the recombinant AAV virion into a suitable muscle cell to produce a transduced muscle cell.

19. The method of claim 15, characterized in that the muscle cell is derived from skeletal muscle.

20. The method of claim 19, characterized in that the muscle cell is a skeletal myoblast or a skeletal myocyte.

21. The method of claim 18, characterized in that the muscle cell is derived from smooth muscle.

22. The method of claim 18, characterized in that the muscle cell is derived from cardiac muscle.

23. The method of claim 22, characterized in that the muscle cell is a cardiomyocyte.



1/9

2/3



FIG. 1



FIG. 2

3/9




FIG. 3

4/9




FIG. 4

5/3

6/9




FIG. 5




FIG. 6

7/9

3/3

# *In vitro* Expression of GAA in Human Skeletal Muscle




FIG. 7





FIG. 8

### Long-term Expression of Human GAA in Mice

9.6



**FIG. 9**

# INTERNATIONAL SEARCH REPORT

INTERNATIONAL APPLICATION: 97/00895

A. CLASSIFICATION OF SUBJECT MATTER	
INC 6	C12H15/18
	C12H15/56
C07K14/505	C12H15/85
A61K31/70	C12H5/10
	C12H9/22
	A61I48/00

RECEIVED BY THE NATIONAL DEFENSE ACADEMY FROM THE NATIONAL DEFENSE ACADEMY  
B. FIELD SEARCHED  
IN THE NATIONAL DEFENSE ACADEMY (classification system followed by classification symbol)  
IRC 6 C129 461K

the effect of the first derivative on the second derivative is investigated in the 3d and 4d models.

הנהגתו של השר לא תהיה כרוכה באחריות. השר יתחבב את המעורבות של השרותים הממשלתיים, וינסה להקטין את המעורבות של השרותים הממשלתיים.

C. DECISIONS CONSIDERED TO BE RELEVANT

	connected with evidence, of the various persons
--	---

**Rekha; 2013a, 50.**

BL00D,  
vol. 86, no. 10, SUPPL. 2, 15 November  
1995.

11-14,  
18-20,  
22, 23

see the whole document

vol. 8, no. 1, January 1995  
page 178 XP000673-01

11, 13,  
28, 22, 23

MOURIS, A. P., ET AL.: "CANDIDATE  
THERAPY WITH ADENO-ASSOCIATED VIRUS AS A  
MEANS OF ACHIEVING GRAFT-SPECIFIC  
IMMUNE-SUPPRESSION".  
see the whole document

1

Special categories of cited documents !

☒ **Alert:** Family members are listed in arrest

considered to be of particular relevance

שנת ה'תשס"ה

which is cited as evidence of another abuse of the special provision

**DECLARATION OF INTEREST**

I, \_\_\_\_\_, declare that I am not a shareholder or owner of any interest in the company.

later gives the priority due credit

examples of the unramified

16 May 1997

Attending Address of the CIA

LE 50001 FIELD: CHINA, F.H. SHI FAYEILIAN ?  
 NL - 2201 HV RIJEPA  
 TEL: 11-70 10045 TO 1158

FILE (031-5) 240206

1 D. 66. 97

4/24/2019 10:00:00

10

Holztorf, 3

Page 1 of 2

International Application No.  
PCT/US 97/06395

## INTERNATIONAL SEARCH REPORT

המחלקה לבריאות הציבור

Control Application No  
PCT/US 97/008895

Classification	Document	Document	Document
1-23	<p>NO 95 J4610 A (CHILDRENS HOSPITAL INC) 21 December 1995 see page 9</p> <p>--- HUMAN GENE THERAPY, vol. 6, October 1995, pages 1329-1341, XP000563710 CLARK, K.R., ET AL. : "CELL LINES FOR THE PRODUCTION OF RECOMBINANT ADEIN-ASSOCIATED VIRUS" see the whole document</p> <p>--- PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 91, November 1994, pages 11557-11561, XP000388922 TRIPATHY, K.S., ET AL. : "STABLE DELIVERY OF PHYSIOLOGIC LEVELS OF RECOMBINANT ERYTHROPOIETIN TO THE SYSTEMIC CIRCULATION BY INTRAVASCULAR INJECTION OF REPLICATION-DEFECTIVE ADEINOVIRUS" see the whole document</p> <p>--- TRENDS IN GENETICS, vol. 9, no. 8, 1993, pages 269-274, XP02030823 BLAU, N.H., ET AL. : "MYOBLASTS IN PATTERN FORMATION AND GENE THERAPY" see the whole document</p> <p>----</p>	1-23	1-23

Parent document and research paper	Publication date	Parent family number(s)	Publication date
NO 93146/0 A	21-12-95	AU 3124395 A CA 2392215 A EP 0764213 A	05-01-96 21-12-95 26-03-97